

A multiscale maximum entropy moment closure for locally regulated space–time point process models of population dynamics

Michael Raghib · Nicholas A. Hill · Ulf Dieckmann

Received: 13 October 2009 / Revised: 29 March 2010 / Published online: 6 May 2010
© Springer-Verlag 2010

Abstract The prevalence of structure in biological populations challenges fundamental assumptions at the heart of continuum models of population dynamics based only on mean densities (local or global). Individual-based models (IBMs) were introduced during the last decade in an attempt to overcome this limitation by following explicitly each individual in the population. Although the IBM approach has been quite useful, the capability to follow each individual usually comes at the expense of analytical tractability, which limits the generality of the statements that can be made. For the specific case of spatial structure in populations of sessile (and identical) organisms, space–time point processes with *local regulation* seem to cover the middle ground between analytical tractability and a higher degree of biological realism. This approach has shown that simplified representations of fecundity, local dispersal and density-dependent mortality weighted by the local competitive environment are sufficient to generate spatial patterns that mimic field observations. Continuum approximations of these stochastic processes try to distill their fundamental properties, and they keep

M. Raghib is grateful for a postgraduate scholarship from the Principal's development fund of the University of Glasgow, an overseas student award granted from the Department of Mathematics, University of Glasgow and DARPA (Award ID:HR001-05-1-0057).

M. Raghib · N. A. Hill
Department of Mathematics, University of Glasgow, Glasgow, UK

Present Address:
M. Raghib (✉)
Program in Applied and Computational Mathematics,
Department of Ecology and Evolutionary Biology,
Princeton University, Princeton, NJ, USA
e-mail: mraghib@princeton.edu

U. Dieckmann
Ecology and Evolution Program, International Institute for Applied Systems Analysis,
Laxenburg, Austria

track of not only mean densities, but also higher order spatial correlations. However, due to the non-linearities involved they result in infinite hierarchies of moment equations. This leads to the problem of finding a ‘moment closure’; that is, an appropriate order of (lower order) truncation, together with a method of expressing the highest order density not explicitly modelled in the truncated hierarchy in terms of the lower order densities. We use the principle of constrained maximum entropy to derive a closure relationship for truncation at second order using normalisation and the product densities of first and second orders as constraints, and apply it to one such hierarchy. The resulting ‘maxent’ closure is similar to the Kirkwood superposition approximation, or ‘power-3’ closure, but it is complemented with previously unknown correction terms that depend mainly on the avoidance function of an associated Poisson point process over the region for which third order correlations are irreducible. This domain of irreducible triplet correlations is found from an integral equation associated with the normalisation constraint. This also serves the purpose of a validation check, since a single, non-trivial domain can only be found if the assumptions of the closure are consistent with the predictions of the hierarchy. Comparisons between simulations of the point process, alternative heuristic closures, and the maxent closure show significant improvements in the ability of the truncated hierarchy to predict equilibrium values for mildly aggregated spatial patterns. However, the maxent closure performs comparatively poorly in segregated ones. Although the closure is applied in the context of point processes, the method does not require fixed locations to be valid, and can in principle be applied to problems where the particles move, provided that their correlation functions are stationary in space and time.

Keywords Birth–dispersal–death stochastic processes · Hierarchies of moment equations · Spatially structured plant population dynamics · Biologically driven spatial pattern formation · Model reduction

Mathematics Subject Classification (2000) 92B99 · 60K35 · 82B31

1 Introduction

One of the most widely used models in theoretical ecology is the logistic equation (Murray 1993; Pearl and Reed 1920; Verhulst 1838)

$$\begin{aligned} \frac{d}{dt}m_1(t) &= r m_1(t) \left(1 - \frac{m_1(t)}{K}\right) \\ m_1(0) &= n_0, \end{aligned} \quad (1)$$

which describes the dynamics of a population in terms of a single state variable $m_1(t)$, which can be interpreted as the expected population size or as the global density. The rate of change of the density in the logistic model is determined by three drivers. The first two are present in the net growth term $r = b - d$, where b and d are respectively the *per capita* fecundity and intrinsic mortality rates. The third one is the density-dependent mortality rate, which is assumed to be proportional to the density, with constant of proportionality $1/K$, where K is the ‘carrying capacity’, i.e. the maximum number of individuals per unit area or volume that can be supported

by some unspecified limiting resource. This model is built on the following set of assumptions (Bolker and Pacala 1997; Dieckmann and Law 2000; Law et al. 2003):

1. There are no facilitative interactions among conspecifics.
2. Contributions to mortality due to competition are pairwise additive.
3. The limiting resource is uniformly distributed in space.
4. There are no differences among individuals in age, size or phenotype.
5. The spatial locations of the individuals are uncorrelated.
6. Allocation to reproduction is independent of the local resource availability.
7. Density-dependent mortality occurs at the same temporal scales than fecundity and intrinsic mortality.

These assumptions are valid only for a rather restricted set of biological situations. For instance, facilitative interactions are known to play a determinant role alongside competition in shaping community structure and dynamics (Brooker et al. 2008). In plant communities, non-successional positive interactions can result from additional resources being made available through synergies (e.g. hydraulic lift, microbial enhancement, mycorrhizal networks), a reduction in the impact of climate extremes and predation (Gross 2008) or a combination of these. The assumption of pairwise additivity in density-dependent mortality enjoys some degree of empirical support for plant populations (Weigelt et al. 2007), but it is still an unresolved issue (Damgaard 2007; Dormann and Roxburgh 2005). Forms of population structure driven by size (or age), phenotype or spatial pre-patterning in the abiotic substrate having an impact on fecundity, recruitment and survivorship are ubiquitously observed both in the field and experimental literature (Purves and Law 2002; Silvertown and Doust 1993; Turnbull et al. 2007). Seed dispersal and competitive interactions are known to occur over a characteristic range of spatial scales rather than being uniformly distributed as is commonly assumed in the logistic model (Condit et al. 2000; Dale 1999; Gratzler et al. 2004; Schneider et al. 2006; Silvertown and Doust 1993; Silvertown and Wilson 2000; Stoll and Weiner 2000).

These limitations have motivated the search for alternatives to the logistic equation that can address questions of broader biological interest, while simultaneously maintaining a reasonable degree of mathematical and computational tractability. Achieving this goal depends heavily on the development of multiscale modeling approaches capable of linking patterns manifested at the larger, population-level scales, to their drivers, which lie in biological processes occurring at the level of individuals. These typically involve spatial and temporal scales that differ substantially from those at which the population-level regularities are detected (Bolker and Pacala 1997; Bolker et al. 2003; Dieckmann and Law 2000; Durrett 1999; Law and Dieckmann 2000; Law et al. 2003; Moorcroft and Pacala 2001; Levin 1994; Satō and Iwasa 2000).

Among all the possible paths suggested as one relaxes these assumptions (1–7), understanding the role of spatial structure, particularly that driven by biological processes alone, has received a considerable amount of interest (Bolker and Pacala 1997, 1999; Bolker et al. 2000; Borgogno et al. 2009; Dale 1999; Dieckmann et al. 2000; Durrett and Levin 1994; Filipe and Gibson 1998, 2001; Filipe and Maule 2003; Filipe et al. 2004; Hiebeler 1997, 2006; Keeling et al. 1997; Keeling 1999; Keeling et al. 2000; Keeling and Ross 2009; Iwasa 2000; Law et al. 2003; Lewis and Pacala

2000; Ovaskainen and Cornell 2006a,b; Pascual and Levin 1999; Pascual et al. 2001, 2002; Picard and Franc 2001; Scanlon et al. 2007). The approaches that have been developed for the spatial problem have a number of commonalities. They usually consist of an individual-based model (IBM) (DeAngelis and Mooij 2005; Grimm 1999) which follows simplified representations of the life histories of each individual in the population. These representations include the biological processes believed to play a role in driving the population-level phenomena, and typically include a combination of fecundity, dispersal, mortality and in some cases, growth. These are modeled in such a way that some form of density-dependent regulation is present in at least one of them. Second, the density-dependent regulation is determined by the neighborhood configuration surrounding each focal individual, which leads to a *local* regulation of the process (Blath et al. 2007; Etheridge 2004; Fournier and Méléard 2004). Third, the dynamics of the macroscopic patterns is obtained from an ensemble average of a sufficiently large number of independent realisations of the individual-level model. Insights about the emergence of various forms of population structure, in particular space, are gained as these broad scale patterns are allowed to vary with the characteristic scales that regulate the biological processes at the level of the individual organism (Bolker and Pacala 1997; Law and Dieckmann 2000; Pacala and Levin 1997; Pascual and Levin 1999; Wilson and Keeling 2000).

This approach, albeit quite illuminating, restricts severely statements that can be made about how the various processes and scales involved interact to produce pattern, since typically there is an absence of a model that summarizing the dynamics of pattern at the larger scales. To circumvent this deficiency, several attempts to derive population-level models from the IBM have been introduced in the literature, for both discrete (Filipe and Gibson 1998, 2001; Filipe and Maule 2003; Hiebeler 2006; Iwasa 2000; Keeling 1999; Ovaskainen and Cornell 2006b) and continuous (Bolker and Pacala 1997, 1999; Dieckmann and Law 2000; Law and Dieckmann 2000; Law et al. 2003; Murrell et al. 2004) representations of space. In the context of spatial pattern in plant population dynamics (Bolker and Pacala 1997; Iwasa 2000; Law and Dieckmann 2000; Scanlon et al. 2007), these models typically take the form of hierarchies of equations for relevant families of summary spatial statistics where quantities in addition to the mean density capture spatial correlations among pairs, triplets etc, that quantify spatial pattern across a range of scales (Dale 1999; Stoyan et al. 1995). These summary statistics are closely related to the central, factorial or raw spatial moments of the underlying spatial stochastic process. For pair configurations in plant population models, common choices are the spatial auto-covariance or the second order product density (Dale 1999; Daley and Vere-Jones 2003; Diggle 1983; Stoyan et al. 1995). A discussion of these various approaches in the development of continuum approximations to spatio-temporal stochastic processes in ecology can be found in a compilation edited by Dieckmann et al. (Dieckmann et al. (2000)).

The non-linearities involved in the density-dependent rules commonly employed in the formulation of spatially explicit IBMs, inevitably result in infinite hierarchies of evolution equations for the summary statistics, where the dynamics of the correlations of order k is tied to that of order $k + 1$. The hierarchy will grow even faster if non-linearities of order higher than quadratic are involved in the dynamics. If one truncates the hierarchy at some order, the evolution equation at the order of the truncation

will depend on the *unknown* density of the next higher order. Analysis of these hierarchies can only proceed after truncation for some small order. This requires the solution of two problems. The first, is identifying an appropriate order of truncation k . The second is compensating for the resulting loss of information. The order of truncation in existing models is chosen on the basis of computational complexity, and rarely goes beyond two (Singer 2004; Bolker and Pacala 1997; Law and Dieckmann 2000). For the second problem, the density of order $k + 1$ is replaced by a functional relationship of all the densities of order up to k , usually called a ‘moment closure’. This functional dependence of higher order quantities on lower order ones is constructed mainly on heuristic reasoning (Bolker and Pacala 1997; Dieckmann and Law 2000; Murrell et al. 2004). For instance, when the order of truncation is two, assuming vanishing central moments of order three leads to the so-called ‘power-1’ closure (Bolker and Pacala 1997). The ‘power-2’ closure arises from an analogy with the pair approximation used in discrete spatial models (Dieckmann and Law 2000; Iwasa 2000). Assuming independence of the three pair correlations associated with each edge of a triplet for all spatial scales leads to the ‘power-3’ or Kirkwood superposition approximation (Dieckmann and Law 2000; Kirkwood and Boggs 1942). More elaborate ‘mixed’ closures involving combinations of these three basic forms (Filipe and Maule 2003), or the introduction of weighting constants (Law et al. 2003; Murrell et al. 2004) have also been proposed. Although closures for orders higher than three do exist, they have restricted applicability due to the daunting computational problem that results (Singer 2004).

Despite some encouraging success that resulted in analytical solutions of the hierarchy at equilibrium for truncation at second order using the ‘power-1’ closure (Bolker and Pacala 1997, 1999; Bolker et al. 2000), remarkably good fit of the numerical solution of the hierarchy with individual-based simulations with so-called *asymmetric* versions of ‘power-2’ closures (Law et al. 2003; Murrell et al. 2004), or the formal derivation of power series expressions for the mean density (Ovaskainen and Cornell 2006a,b), most predict poorly the equilibrium densities even for situations of mild spatial correlations. In the cases where they succeed over broader ranges of spatial correlations (i.e. the asymmetric power-2), the closure depends on tuning a set of weighting constants whose values can presently be found only by comparison with simulations of the stochastic process, and its success is entirely model-specific (Ovaskainen and Cornell 2006b). A significant obstacle in the widespread adoption of these continuum approximations and their closures is that none is equipped with a criterion for their domain of validity beyond comparisons with simulations of the individual-based model, an exception being the work of Ovaskainen and Cornell (Ovaskainen and Cornell 2006a,b) who provided error bounds, but require long range—albeit finite—interactions to provide accurate results. Nevertheless, many of these heuristic closures do provide a better approximation to the dynamics of a spatially structured population than the logistic equation, and illuminate a variety of mechanisms by which endogenously generated spatial pattern appears in plant populations.

Inspired by earlier results of Hillen (2004) and Singer (2004), who used the principle of constrained maximum entropy (Khinchin 1957; Jaynes 1957; Shannon and Weaver 1949) to respectively derive closures for velocity jump processes (Othmer et al. 1988) and the BBGKY hierarchy arising in the statistical mechanics of fluids

(Kirkwood and Boggs 1942), we develop a closure scheme based on constrained entropy maximisation for the moment hierarchy developed by Law and Dieckmann (2000), constrained to satisfy normalisation and the product densities up to order two. In order to be able to relate the output of the entropy maximisation to the approximating dynamical system, we also reframe the hierarchy of Law and Dieckmann (2000) in terms of product densities rather than the spatial moments. These two kinds of sets of summary statistics are very closely related, since the latter can be seen as estimators of the former. The approach of Hillen (2004) consists of proving that the L^2 -norm over the space of velocities of the transport equation of Othmer et al. (1988) behaves like an entropy, with the velocity moments acting as constraints. Singer (2004) treats the triplet product density as a probability density in order to construct an entropy from the point of view of information theory (Jaynes 1957; Khinchin 1957; Shannon and Weaver 1949), using consistency of the marginals as constraints. Our approach differs from these two maximum entropy maximisation methods in a number of ways. First, we use the information theoretical entropy functional for point processes (Daley and Vere-Jones 2004; McFadden 1965), defined by the negative of the expected log-likelihood. This definition includes all the orders that contribute spatial information. Second, the product densities, which come from the moment hierarchy and provide the constraints, are incorporated into the entropy functional by means of an expansion that allows to express the likelihoods (or Janossy densities) in terms of product densities and vice versa (Daley and Vere-Jones 1988, 2003). This allows to establish a formal connection between the entropy functional and the moment hierarchy. Third, our closure is implicit, in the sense that the density of order three appears on both sides of the closing relationship, thus allowing irreducible correlations of third order to be explicitly included. Fourth, the method presented here complements the Kirkwood (or power-3) closure with previously unknown correction terms that depend on the area for which the three points in the triplet become independent. These correction terms are important where the three particles in the triplet configuration are close to each other, but progressively vanish as these become separated, at which point the maximum entropy closure reduces to the classical Kirkwood superposition approximation. In addition, the closure comes equipped with a criterion of validity stemming from the normalisation constraint. This validity check comes from an ancillary integral equation that returns the area of the domain for which a third point in the triplet becomes independent of the other two. This equation produces a single, non-trivial root when the correlations predicted by the moment hierarchy are consistent with the truncation assumptions, but fails to do so otherwise. The maximum entropy closure relationship we found is given by

$$m_3(\xi_1, \xi_2) = \left[m_2(\xi_1) - |A_0| \int_{A_0} m_3(\xi_1, \xi'_2) d\xi'_2 \right] \\ \times \left[m_2(\xi_2) - |A_0| \int_{A_0} m_3(\xi_2, \xi_2 - \xi'_1) d\xi'_1 \right]$$

$$\begin{aligned} & \times \left[m_2(\xi_2 - \xi_1) - |A_0| \int_{A_0} m_3(\xi_2 - \xi'_1, \xi'_1) d\xi'_1 \right] \\ & \times \frac{\exp(-m_1|A_0|)}{\left[m_1 - |A_0| \int_{A_0} m_2(\xi'_1) d\xi'_1 + \frac{|A_0|^2}{2} \int_{A_0 \times A_0} m_3(\xi'_1, \xi'_2) d\xi'_1 d\xi'_2 \right]^3}, \end{aligned} \tag{2}$$

where m_1, m_2, m_3 are the first, second and third order product densities (the densities of the factorial moments of the underlying spatial point process), ξ_1 , and ξ_2 are vector distances respectively linking the pairs of particles (x_1, x_2) and (x_1, x_3) forming a triplet configuration. The set A_0 is a circular domain of area $|A_0|$ that establishes the spatial scale within which triplet correlations are irreducible. The exponential term is the avoidance function (i.e. the probability of observing no points in A) of an ancillary Poisson point process on the window A with the same mean density m_1 as the correlated process that generates the hierarchy (Daley and Vere-Jones 1988, 2003). For this specific window the avoidance function of the correlated process (in general consisting of an infinite series in terms of product densities of all orders rather than an exponential function) coincides with that of the ancillary Poisson point process, which is exponential. This set is found as the domain of integration that solves the normalisation condition

$$\int_{A_\epsilon} m_2(\xi'_1) d\xi'_1 - \frac{1}{3} \int_{A_\epsilon \times A_\epsilon} m_3(\xi'_1, \xi'_2) d\xi'_1 d\xi'_2 = |A_\epsilon| m_1^2 - \frac{|A_\epsilon|^2}{3} m_1^3 \tag{3}$$

where A_ϵ is a circular domain of radius ϵ centered at the origin. The set A_0 is found by allowing the radius ϵ to take positive real values until the equality in (3) holds. This closure is applied if the three points in the triplet lie inside A_0 , and outside this region the classical Kirkwood closure applies. The largest correction by far is due to the exponential term, since the integral correction terms in the numerator and denominator tend to be of similar magnitude, in which case the maxent closure is simply given by

$$m_3(\xi_1, \xi_2) \approx \frac{m_2(\xi_1) m_2(\xi_2) m_2(\xi_2 - \xi_1)}{m_1^3} \exp(-m_1|A_0|), \tag{4}$$

where the vector distances ξ_1, ξ_2 and $\xi_2 - \xi_1$ that connect the three points in the tripled fall outside the domain A_0 , the closure reduces to the classical Kirkwood superposition approximation or ‘power-3’ closure, which can be justified using probabilistic independence arguments (Kirkwood and Boggs 1942; Singer 2004), given by

$$m_3(\xi_1, \xi_2) = \frac{m_2(\xi_1) m_2(\xi_2) m_2(\xi_2 - \xi_1)}{m_1^3}. \tag{5}$$

The normalisation constraint (3) acts as a criterion of validity in the sense that it will only find a single nontrivial domain A_0 when the assumptions of the closure are

met. A set of measure zero is always a solution for (3), thus failure to find a single additional non-trivial solution can occur when:

1. The radius ϵ grows without bound as one iterates (3) and (4) to find A_0 . This implies that higher order terms are required for an accurate representation of the triplet density or that spatial correlations decay slowly.
2. There is a finite number of solutions to (3) which implies that there are multiple scales of spatial pattern (i.e. segregated clusters).

We find that the correction terms in the maxent closure lead to substantial improvements in the prediction of the equilibrium density for mildly aggregated patterns when compared with heuristic closures. For highly clustered patterns that involve different types of pattern over a range of scales, for instance a combination of aggregation at short scales and segregation at intermediate ones, the method fails to find a non-trivial root for the area A_0 involved in the correction terms, which indicates that the combination of parameters that lead to this correlation regime is outside the domain of validity. Somewhat surprisingly the Kirkwood closure behaves better than the maxent closure 2 for segregated (or regular patterns). This occurs because in this regime short range segregation at very short scales is compensated by clustering at intermediate scales, and the ancillary equation for A_0 detects only the aggregation scale, and not the segregation one, which leads to over correction.

The paper is organized as follows. Section 2 discusses the locally regulated space-time point process model originally developed by Law and Dieckmann (2000), and includes a Gillespie-type simulation algorithm (Gillespie 1976; Renshaw 1991), together with known definitions and estimators for the product densities and some simulation results included for illustration purposes. Simulation results exploring a broader range of parameters for this process can be found elsewhere (Bolker et al. 2000; Law et al. 2003; Murrell et al. 2004; Raghیب Moreno 2006). Section 3 reframes the spatial moment equations of Law et al. Law et al. (2003) in terms of product densities. Section 4 discusses the moment closure for truncation at second order based on constrained entropy maximisation. Section 5 discusses the numerical implementation of the closure and compares its predictions against simulations of the point process for mildly aggregated patterns. Finally, Sect. 6 presents a critique of the maximum entropy method, and suggests further areas of development.

2 Spatio-temporal point process model

We consider a single population of identical individuals, each of which can occupy arbitrary locations on a 2-dimensional torus A . The state of the population for each fixed time t is modeled as a realisation of a spatial point process, called the configuration or point pattern (Daley and Vere-Jones 2003; Diggle 1983; Stoyan et al. 1995),

$$\varphi_t(A) = \{x_1, \dots, x_{N_t}\}, \quad (6)$$

where the x_i are the spatial locations of all individuals found within A . Alternatively we have

$$N_t(A) = \# \{x_1, \dots, x_{N_t}\}$$

where $N_t(A)$ stands for the total population counts within A , and the cardinality operator $\#$ counts the number of elements in a set. Note that in (6) both the locations x_i and the total counts N_t are random variables. The dynamics of the population is modeled by introducing a time component, where the updating times are also random variables, subject to *local* regulation (Cressie 1991; Daley and Vere-Jones 2003). Two versions of this model have been introduced independently by Bolker and Pacala (1997) and Dieckmann and Law (2000). Both share the key ingredients of non-uniform dispersal, and a density-dependent mortality term that depends on the configuration surrounding the focal individual which is the mechanism that introduces the local regulation. The configuration (6) evolves in time by sampling from two exponential distributions of waiting times that regulate the inter-event times between fecundity/dispersal and mortality events at the individual level, where the latter is determined from both intrinsic and density-dependent contributions.

2.1 Dispersal and fecundity

Per capita waiting times between births are assumed to be exponentially distributed with constant parameter b , the *birth (or fecundity) rate*. If a birth occurs, the newborn is displaced instantaneously from the location of its mother x_i to a random new location x_j , sampled from the probability density, $B(x_i - x_j; \sigma_B)$ the *dispersal kernel*, where σ_B is a parameter that measures the characteristic dispersal length. The index i of the mother is chosen uniformly from the list of indices $J_A = \{1, 2, \dots, N_t(A)\}$ in the configuration.

2.2 Mortality

The probability that a given individual i at location x_i dies in the time interval $(t, t + dt)$ is also assumed to be exponentially distributed with parameter $m(x_i)$, the *total per capita mortality rate*, given by

$$m(x_i) = d + d_N \sum_{j \neq i \in J_A} W(|x_i - x_j|; \sigma_W), \tag{7}$$

where d , is the *intrinsic* mortality rate, and d_N is the *density-dependent* mortality rate. In order to allow comparisons with the predictions of the logistic model (1) we defined it as $d_N = (b - d)/K$, where K is the non-spatial carrying capacity (the expected value at equilibrium under complete spatial randomness). This second ‘mortality clock’ is rescaled by a weighted average of the local configuration around the focal individual, so that mortality due to competition is more likely to occur in locally dense regions than in comparatively sparse ones. The contributions of neighbors to

Table 1 Point process model parameters

Parameter	Symbol	Units
Fecundity	b	time^{-1}
Intrinsic mortality	d	time^{-1}
Density-dependent mortality	d_N	$\text{time}^{-1} \text{ indiv}^{-1}$
Non-spatial carrying capacity	K	individuals
Dispersal scale	σ_B	length
Competition scale	σ_W	length
Initial population size	N_0	individuals
Spatial arena	A	length^2

the mortality of x_i are assumed to decay monotonically with distance. This is modeled by a normalized, radially symmetric weighting function $W(|\xi|; \sigma_W)$, the *mortality kernel*, that vanishes outside a finite interaction domain D_W , where σ_W is a parameter associated with the characteristic length scale of competitive interactions. This function is interpreted as an average effect that simplifies the details of the physiology of mortality due to crowding. The parameters of the model are summarized in Table 1.

2.3 Simulation algorithm

A sample path for the space–time point process with rates described in Sects. 2.1 and 2.2 can be simulated by a variant of the Gillespie algorithm (Gillespie 1976; Renshaw 1991). The spatial arena can be identified with the unit square $W = [0, 1] \times [0, 1]$ (after rescaling the parameters in the interaction kernels), with periodic boundary conditions. The initial population consists of N_0 individuals, and $[0, T_{\max}]$ is the time interval of interest.

1. Generate the configuration at time $t = 0$, $\varphi_0 = \{(x_1, y_1); \dots; (x_{N_0}, y_{N_0})\}$, from two independent sets of N_0 deviates from $U(0, 1)$, $X_0 = \{x_1, \dots, x_{N_0}\}$ and $Y_0 = \{y_1, \dots, y_{N_0}\}$.
2. While the elapsed time t is less than T_{\max} do:
 - (a) Generate a birth waiting time T_b from the exponential density with parameter $b N_t$, where N_t is the number of individuals that are alive at time t .
 - (b) Generate the set of mortality waiting times $T_m = \{\tau_1, \dots, \tau_{N_t}\}$ from a set of N_t exponential random variables, each with parameter $m(x_i) = d + d_N \sum_{j \neq i} W(|x_i - x_j|)$, for each of the $i = 1, \dots, N_t$ individuals in the configuration at time t
 - (c) The time until the next event is given by $\tau_n = \min\{T_b \cup T_m\}$.
 - i. A birth occurs if $\tau_n = T_b$, in which case the location of the newborn individual x_b is given by

$$x_b = x_p + \xi$$

where the index of the parent p is drawn uniformly from the set of indices $I_\varphi(t) = (1, \dots, N_t)$ and the displacement ξ is drawn from the

dispersal kernel $B(\xi)$. The configuration is then updated to include the newborn

$$\varphi_{t+T_b} \rightarrow \varphi_t \cup \{x_b\}.$$

- ii. If $\tau_n \neq T_b$ then the next event is a death in which case the i th individual in T_m for which $\tau_i = \tau_n$ is removed from the configuration

$$\varphi_{t+\tau_n} \rightarrow \varphi_t \setminus \{x_i\}$$

- (d) Update the elapsed time $t \rightarrow t + \tau_n$.

2.4 Summary statistics

The specific configurations resulting from simulations of the algorithm in Sect. 2.3 are of limited interest. The fundamental question is understanding how spatial correlations develop from an unstructured initial condition, and how the equilibrium density departs from the logistic behavior when considering an ensemble of simulations for various combinations of the spatial scales of competition and dispersal (Bolker and Pacala 1997; Dieckmann and Law 2000; Law and Dieckmann 2000). This requires a set of summary statistics capable of distinguishing various forms of spatial structure for the same population size. This is illustrated in the upper three panels in Fig. 1) that shows three types of point configurations with the same number of points. A useful set for this task are the product densities (or densities of the factorial moments), i.e. the densities of the expected configurations involving one, two or more *distinct* points after removing self-configurations (Daley and Vere-Jones 2003; Diggle 1983; Stoyan et al. 1995). For spatially stationary point processes, these are functions of the inter-point distances between the points comprising an expected configuration of a certain order k . The product densities are defined in terms of the population count $N_t(B)$ observed through some window B at time t defined as (Cressie 1991; Daley and Vere-Jones 2003; Stoyan et al. 1995)

$$N_t(B) = \sum_{x_i \in \varphi_t} I_B(x_i), \tag{8}$$

where $I_B(x)$ is the indicator function of the set B defined by

$$I_B(x) = \begin{cases} 1 & \text{if } x \in B, \\ 0 & \text{otherwise.} \end{cases} \tag{9}$$

The coarsest is the mean density (or intensity) which measures the expected number of individuals per unit area at each time, defined as

$$m_1(x, t) = \lim_{\epsilon \downarrow 0} \frac{E\{N_t(S_\epsilon(x))\}}{|S_\epsilon(x)|} \tag{10}$$




Fig. 1 The three upper panels show different types of point patterns sharing the same number of points $N(A) = 500$, where the window A is the unit square. **a** Shows aggregation at short scales, but segregation at intermediate ones. **b** Corresponds to complete spatial randomness and **c** displays a segregated (or regular) pattern. In **a**, there is a tendency of points to occur near each other and form clusters, but the clusters themselves tend to be segregated. It is difficult to distinguish **b** and **c** visually but points tend to avoid each other over short scales in **c**, whereas some clusters can be observed in **b**. The three lower panels show estimates of the pair correlation function (15) for each of the point patterns above. The dashed lines in the lower panels correspond to the 99% Monte Carlo envelopes

where $S_\epsilon(x)$ is the open ball of radius ϵ centered around x , and $|A|$ is the area of the window A . Since the mortality and fecundity rates do not depend specific locations but on relative distances, and both the dispersal and competition kernels are symmetric by definition, the spatial point process is spatially stationary and isotropic, in which case the mean density is constant for each fixed time

$$m_1(x, t) = m_1(t).$$

A naïve estimator for the mean density from a single realisation is (Diggle 1983; Stoyan et al. 1995)

$$\hat{m}_1(t) = \frac{N_t(A)}{|A|} \tag{11}$$

where $N_t(A)$ is as in (8). If an ensemble of Ω independent replicates of the process is available, this estimate can be improved by averaging over the ensemble

$$\bar{m}_1(t) = \frac{\langle N_t(A) \rangle_\Omega}{|A|}. \tag{12}$$

For a Poisson process, the mean density (10) is a sufficient statistic for the process. More general cases require keeping track of spatial correlations. Higher order quantities are required to distinguish between aggregated (or clustered), random and segregated (or over-dispersed) point patterns with the same mean density (see Fig. 1). For this purpose we need, at the very least, information about two-point correlations. These are measured by the pair correlation function, defined as the ratio

$$g_2(\xi; t) = \frac{m_2(\xi, t)}{m_1^2(t)} \tag{13}$$

which requires knowledge of the density of the expected number of pairs at spatial lag ξ , measured by the second order product density $m_2(\xi, t)$.

$$m_2(\xi; t) = \lim_{\epsilon \downarrow 0} \frac{\mathbb{E} \{ N_t(S_\epsilon(\mathbf{0})) [N_t(S_\epsilon(\mathbf{0} + \xi)) - \delta_{\mathbf{0}}(S_\epsilon(\mathbf{0} + \xi))] \}}{|S_\epsilon(\mathbf{0})| |S_\epsilon(\mathbf{0} + \xi)|} \tag{14}$$

where $S_\epsilon(\mathbf{0})$ and $S_\epsilon(\mathbf{0} + \xi)$ are small windows of observation respectively centered at the origin, and at distance ξ from the origin. The Dirac measure in the second factor in the numerator removes the count at zero lag from the second window in order to avoid self-configurations. In general, the definition (14) centers the count for each specific location x , but given that in our case the process is stationary and isotropic by construction, it can be translated to the origin without loss of generality, in which case m_2 depends only on the spatial lag ξ .

In the case of a spatially random configuration (a Poisson point process), the counts on non-overlapping windows are independent of each other and thus the second order density is simply the square of the mean density. Correlations of configurations involving k points are simply the k th powers of the mean density (Diggle 1983; Stoyan et al. 1995). The pair correlation function (13) is the lowest order product density that allows detection of departures from complete spatial randomness. Thus, values of the pair correlation function greater than one for some lag ξ indicate aggregation at that scale, whereas values below one signal segregation. Estimation of the pair correlation function requires an estimator of the squared density (Stoyan et al. 1995)

$$\bar{m}_1^2(t) = \frac{\langle N_t(A) [N_t(A) - 1] \rangle_\Omega}{|A|^2},$$

together with a kernel density estimator for the second order product density (Scott 1992; Stoyan and Helga 1994),

$$\hat{m}_2^{(h)}(r, t) = \frac{1}{2\pi r} \sum_i \sum_{j \neq i} \frac{k_h(r - \|x_i - x_j\|)}{|A_{x_i} \cap A_{x_j}|} \tag{15}$$

where r is the spatial lag, h is the bandwidth of the kernel density estimate k_h , the points x_i belong to a configuration $\varphi_t(A)$ sampled at time t , and $\|x_i - x_j\|$ is the

Euclidean distance between the points x_i and x_j . The denominator is an edge corrector that rescales the count in the numerator by the area of the intersection of the window of observation A_{x_i} shifted so that its centered around the point x_i , with the window A_{x_j} shifted around x_j (Cressie 1991; Dale 1999; Stoyan et al. 1995)

$$A_{x_i} = \{x + x_i : x \in A\}.$$

If an ensemble of independent realisations is available, the single realisation estimator (15) can be improved by means of an ensemble average

$$\bar{m}_2^{(h)}(r, t) = \left\langle \widehat{m}_2^{(h)}(r, t) \right\rangle_{\Omega}.$$

As before, the angle brackets $\langle \cdot \rangle_{\Omega}$ represent an average of the estimates across a number of independent sample paths Ω . For the smoothing kernel k_h a common choice is the Epanechnikov kernel (Stoyan et al. 1995)

$$k_h(s) = \frac{3}{4h} \left(1 - \frac{s^2}{h^2} \right) I_{(-h,h)}(s),$$

where I is the indicator function (9). Although empirical methods for selection of the bandwidth h are widely used, for instance the rule (Stoyan and Helga 1994)

$$h = c/\sqrt{\widehat{m}_1(t)}, \quad c \in (0.1, 0.2),$$

data-driven methods for optimal choices of h based on cross-validation have been recently introduced (Guan 2007a,b). In general, the product density of order k is defined as (Błaszczyszyn 1995)

$$m_k(x_1, \dots, x_k, t) = \lim_{\epsilon \downarrow 0} E \left\{ \prod_{j=1}^k \frac{[N_t(S_{\epsilon}(x_j)) - \sum_{i=1}^{j-1} \delta_{x_i}(S_{\epsilon}(x_j))]}{|S_{\epsilon}(x_j)|} \right\}, \quad (16)$$

where $\sum_{i=1}^{j-1} \delta_{x_i}(S_{\epsilon}(x_j))$ removes self j -tuples for $j > i$. In the case of spatial stationarity and isotropy, the specific locations x_1, \dots, x_k can be replaced by the relative distances ξ_1, \dots, ξ_{k-1} ,

$$m_k(\xi_1, \dots, \xi_{k-1}, t),$$

and the k th correlation function becomes,

$$g_k(\xi_1, \dots, \xi_{k-1}; t) = \frac{m_k(\xi_1, \dots, \xi_{k-1}, t)}{m_1^k(t)} \quad (17)$$

which is interpreted in a similar way to the pair correlation function, but considering k -plets instead of pairs. The lower three panels in Fig. 1 show the application of second

order summary statistics, particularly the pair correlation function (15) to three types of spatial patterns. Figure 1d indicates at least two scales of pattern. There is short scale aggregation where $\hat{g}_2(r)$ exceeds one, but there is also intermediate scale segregation (or regularity) that indicates that the clusters themselves tend to avoid each other. This is measured in the region where $\hat{g}_2(r)$ is below one. Figure 1e indicates complete spatial randomness, since the pair correlation function is close to one for all values of r . Figure 1c indicates segregation at short scales, since $\hat{g}_2(r)$ falls between 0 and 1 for small values of r . In all the lower panels the dashed lines correspond to the 99% Montecarlo envelopes (Cressie 1991). The insets show that in Fig. 1e, the pair correlation function is confined within the Montecarlo bounds, but in Fig. 1f falls out of the envelopes at short scales.

2.5 Point process simulation results

For the convenience of the reader, simulation results for the point process are shown in Fig. 2, with the same parameter values as in Law et al. (2003), but obtained from code developed independently. The spatial arena is the unit square, and the kernels are both radially symmetric Gaussians, but the mortality kernel is truncated (and renormalized) at $3\sigma_W$. The left panel shows estimates of the mean density versus time for various values of the characteristic spatial scales of dispersal and mortality. The right panel shows the pair correlation function at the end of the simulation for each of the four spatial regimes for which the population persists. Both quantities were estimated from an ensemble of 300 independent sample paths.

Case (a) shows results for a segregated (or regular) spatial pattern that arises from very local competitive interactions, but long range scales of dispersal ($\sigma_B = 0.12$, $\sigma_W = 0.02$). In this situation local densities experienced by a focal individual are lower in average than the random case (the pair correlation function is below one), which results in equilibrium densities that equilibrate at higher values than the non-spatial carrying capacity. This results from the ability of newborns to escape locally crowded regions via the long range dispersal kernel. Case (b) corresponds to dispersal and mortality kernels with large characteristic spatial scales ($\sigma_B = 0.12$, $\sigma_W = 0.12$). In this situation there is enough mixing to destroy spatial correlations—confirmed by its almost constant pair correlation function shown in the right panel—and the mean density equilibrates at a value that is very close to the non-spatial carrying capacity ($K = 200$). Case (c) is associated to a segregated pattern of clusters, which is the converse situation of the segregated pattern with very localized dispersal, and mild competition distributed over a longer range ($\sigma_B = 0.02$, $\sigma_W = 0.12$). The oscillations of the pair correlation function indicate two scales of pattern. There is short scale aggregation, but the clusters themselves form a segregated pattern with respect to each other, so the local crowding due to clustering that should lead to high density-dependent mortality is compensated by the overdispersion. Overall, the local competitive neighborhood experienced by an individual in this situation is more crowded than in a random distribution of points, which results in a mean density that equilibrates at lower values than the non-spatial carrying capacity. Case (d) corresponds to a mildly aggregated pattern ($\sigma_B = 0.04$, $\sigma_W = 0.04$), where




Fig. 2 The left panel shows estimates for the mean density $\bar{n}_1(t)$ from an ensemble of $\Omega = 300$ realisations, and various characteristic spatial scales of dispersal and density-dependent mortality that cover four main types of spatial pattern. The dashed lines are the envelopes for one standard deviation. Kernel parameters are (a) $\sigma_B = 0.12$, $\sigma_W = 0.02$, (b) $\sigma_B = 0.12$, $\sigma_W = 0.12$, (c) $\sigma_B = 0.02$, $\sigma_W = 0.12$, (d) $\sigma_B = 0.04$, $\sigma_W = 0.04$ and (e) $\sigma_B = 0.02$, $\sigma_W = 0.02$. The other parameters, $b = 0.4$, $d = 0.2$, $K = 200$, are fixed for all cases. The spatial arena is the unit square with periodic boundaries. The right panel shows the corresponding estimates for the pair correlation function $\bar{g}_2^*(r)$ at the end of the simulation averaged over the ensemble of simulation runs

there is a single scale of aggregation. Even for small departures from complete spatial randomness such as this one, the effect of the spatial pattern in the dynamics of the mean density is substantial, since we see a reduction of about 30% in the equilibrium density in this case with respect to that of complete spatial randomness. Finally, case (e) indicates an extreme case of aggregation, with very intense, local mortality and dispersal ($\sigma_B = 0.02$, $\sigma_W = 0.02$), where the population goes to extinction (exponentially) after a short growth transient.

3 Moment equations and the closure problem

The central problem associated with the space–time point process described earlier in Sect. 2.3 is to obtain a closed form expression for the finite dimensional distributions,

$$\mathbb{P}_k \{A_1, \dots, A_k, n_1, \dots, n_k; t\}, \tag{18}$$

that determine the probability of observing n_1 points in the window A_1 , n_2 points in the window A_2 , and so forth up to the n_k points in A_k at time t , from the definition of the space–time point process discussed in the previous section. Unfortunately, this seems to be remarkably difficult, due to the presence of the non-linearity in the mortality rate

in (7), and the localized nature of dispersal (Etheridge 2004). However, the question of ecological interest is understanding the modifications that should be introduced to the logistic equation (1) in order to account for the effects of spatial correlations in the dynamics of the mean density. This can be accomplished by deriving evolution equations for the product densities [which are the densities of the factorial moments of (18)] from the transition rates of the point process discussed in the previous section. Following a Master equation approach similar to that used by Bolker and Pacala (1997) and Dieckmann and Law (2000), we derive the following hierarchy of product density equations (see ‘‘Appendix’’). The first member in this hierarchy corresponds to the modified or ‘spatial’ logistic equation (Murray 1992),

$$\frac{d}{dt}m_1(t) = r m_1(t) - d_N \int_{\mathbb{R}^2} W(\xi_1) m_2(\xi'_1, t) d\xi'_1. \tag{19}$$

where $r = b - d$, $d_N = (b - d)/K_s$ and $W(\xi_1)$ is the mortality kernel in (7). K_s is the *spatial* carrying capacity, or the number of individuals per unit area that can be supported under random mixing


$$K_s = \frac{K}{|A|}.$$

Equation (19) shows that the required modification of the logistic equation consists of substituting the quadratic term with an average of the second order product density $m_2(\xi_1, t)$ weighted by the mortality kernel $W(\xi_1)$. This term computes the effective number of neighbors n_{eff} that contribute to density-dependent mortality,

$$n_{\text{eff}}(t) = \int_{\mathbb{R}^2} W(\xi_1) m_2(\xi'_1, t) d\xi'_1.$$

Thus, the effect of mortality on the evolution of the mean density is tied to a weighted average of the mortality kernel with the two-point spatial correlations in the process. Equation (19) reduces to the logistic equation for the Poisson point process, in which case $m_2(\xi_1) = m_1^2$. In aggregated spatial patterns, m_2 exceeds m_1^2 for some domain. If mortality is modeled by a kernel that penalizes close proximity over the same range of scales where aggregation occurs, then the effect of mortality due to competition is stronger in this case than that of the logistic equation, in which case the density equilibrates below K_s [Fig. 2, cases (c)–(e)]. The opposite situation occurs in segregated patterns, where m_2 is less than m_1^2 at the scales where the mortality kernel penalizes aggregation. As a result, the effect of competition on mortality is milder than in a random spatial pattern, in which case the mean density equilibrates at values greater than K_s [Fig. 2, case (a)]. Equation (19) depends on the unknown second order density m_2 . A similar procedure to that used in the derivation of (19) one obtains the evolution equation for this quantity

Fig. 3 Schematic representation of a spatially stationary triplet configuration. The pair densities are evaluated at each inter-event (vectorial) distances ξ_1, ξ_2 and $\xi_1 - \xi_2$



$$\begin{aligned} \frac{1}{2} \frac{d}{dt} m_2(\xi_1, t) = & b \int_{\mathbb{R}^2} B(\xi_2) m_2(\xi_1 - \xi_2, t) d\xi_2 + b B(\xi_1) m_1(t) - d m_2(\xi_1, t) \\ & - d_N W(\xi_1) m_2(\xi_1, t) - d_N \int_{\mathbb{R}^2} W(\xi_2) m_3(\xi_1, \xi_2, t) d\xi_2. \end{aligned} \quad (20)$$

Here the role of the dispersal and competition kernels as the main pattern drivers can be clearly discerned (Bolker and Pacala 1997; Bolker et al. 2000; Dieckmann and Law 2000; Law et al. 2003). The first two terms in (20), related to fecundity and dispersal, are

$$b \int_{\mathbb{R}^2} B(\xi_2) m_2(\xi_1 - \xi_2; t) d\xi_2 + b B(\xi_1) m_1(t).$$

Both are nonnegative by definition for all values of ξ_1 and t . The rate of change of m_2 increases due to their combined effect, and thus they drive aggregation at the scales controlled by the characteristic spatial scale of the dispersal kernel. The convolution measures the creation of pairs along ξ_1 due to dispersal of offspring events generated by the third member of the triplet along the $\xi_1 - \xi_2$ edge (Fig. 3). The second term measures the creation of pairs along the ξ_1 edge due to the dispersal events of offspring generated by the individual at the origin of ξ_1 . The remaining terms, due to mortality are,

$$-d m_2(\xi_1, t) - d_N W(\xi_1) m_2(\xi_1, t) - d_N \int_{\mathbb{R}^2} W(\xi_2) m_3(\xi_1, \xi_2; t) d\xi_2.$$

All the three terms are negative, and thus contribute to the destruction of pairs along the ξ_1 edge, leading to segregated patterns. The first term measures intrinsic mortality of both members of the pair and the remaining ones are related to density-dependent mortality. The second measures mortality of the pairs due to competition at the scales controlled by the mortality kernel. The last term measures the destruction of the pair along the ξ_1 edge due to the effect of competition with the additional member of the triplet located along the ξ_2 edge.

These terms for both dispersal and mortality are initially calculated by fixing the count at the origin of ξ_1 and letting the count at the end of ξ_1 vary according to the fecundity, dispersal and mortality terms. Symmetry considerations require consideration of the reverse situation, where the count at the end of ξ_1 is fixed, and the origin is allowed to vary. Since these are symmetric, these additional terms lead to the factor of $1/2$ on the left hand side of the equation for the second order product density.

4 Moment closure by Shannon entropy maximisation

The product density equations (19) and (20) cannot be solved in that form because the evolution equation for the second order density has a mortality term that depends on a weighted average of the third order one. Although it is possible to derive an additional evolution equation for this quantity, it will involve an unknown fourth order term, leading to a system that is not closed. In general, the evolution equation for the density of order k will depend on the density of order $k + 1$. This gives rise to two problems, known together as ‘a moment closure’ (Bolker and Pacala 1997; Law and Dieckmann 2000). The first is choosing an appropriate order of truncation k , and the second is finding an expression for the product density of order $k + 1$ in terms of the densities of orders up to k (or $k + 1$ in the case of an implicit closure).

Ideally, the order of the truncation should be based on an understanding of the convergence properties of the hierarchy in order to establish error bounds. In practice, the order of the truncation is determined by the computational cost of the numerical solution, which is determined by the size of the arrays that can be stored and operated on efficiently. Explicit representation of third order terms already requires least 3.2 GB of memory using double precision and a relatively coarse discretisation of 100 grid points per dimension. If memory occupies 32 bits per grid point (as is the case in MATLAB), and the two linearly independent vectors ξ_1 and ξ_2 in the triplet density m_3 are 2-d, each requires a 100×100 array to store it. This situation pretty much constrains to three the highest order density that can be represented explicitly.

From an applied perspective, the first and second order terms are of greatest interest, since these respectively encode the dynamics of the average density and the spatial covariance. The latter can be interpreted biologically as the average environment experienced by an individual as a function of spatial scale (Bolker and Pacala 1997, 1999; Bolker et al. 2000; Law and Dieckmann 2000; Law et al. 2003). The shape of the second order correlation function can be used to distinguish between aggregated, random and segregated spatial patterns sharing the same average density (see Sect. 2.4).

Closure problems are pervasive in the statistical mechanics of fluids where thermodynamic quantities are derived from the statistical properties of the particle distributions (Grouba et al. 2004; Kirkwood and Boggs 1942; Meeron 1957; Salpeter 1958; Sese 2005; Singer 2004). Here our intent is somewhat similar in the sense that a detailed individual-based model is used to inform a mean-field model that does not neglect the role of spatial fluctuations in density due to endogenously generated spatial structure (Bolker and Pacala 1997, 1999; Law et al. 2003). Within spatial ecology, moment closures have been proposed with varying degrees of success, using a suite of methods, among which we have:

- *Heuristic reasoning*, where consistency arguments are used to construct closing relationships (Bolker and Pacala 1997; Dieckmann and Law 2000; Law et al. 2003; Murrell et al. 2004).
- *Distributional properties*, where closures are based on assuming a functional form for the distribution of the process (Krishnarajah et al. 2005).
- *Variational methods*, where it is assumed that the unknown distribution optimizes some meaningful functional, usually an entropy-like object (Hillen 2004; Singer 2004)

In order to make the paper reasonably self-contained, we shall briefly review closures based on heuristic reasoning, which have dominated work in this problem. Additional information can be found in a recent review by Murrell et al. (2004).

4.1 Heuristic methods of moment closure

Heuristic closures are usually based on self-consistency arguments. For instance, they should be strictly positive and invariant under permutations of the arguments (Cressie 1991; Daley and Vere-Jones 2003; Diggle 1983). Also, if correlations are assumed to decay monotonically with distance, then there is a distance d beyond which the particles become uncorrelated and thus higher order densities become simple powers of the mean density. Although a large number of functional forms can be chosen in order to satisfy these minimum requirements, the simplest ones usually involve additive combinations of various powers of the second and first moments. For instance, if one further assumes that central third moments vanish, the resulting expansion in terms of product densities, leads to the *power-1* closure, dubbed that way because the highest occurring power of the second order density is one (Bolker and Pacala 1997, 1999; Bolker et al. 2000; Dieckmann and Law 2000),

$$m_3(\xi_1, \xi_2) = m_1 m_2(\xi_1) + m_1 m_2(\xi_2) + m_1 m_2(\xi_1 - \xi_2) - 2 m_1^3. \quad (21)$$

This closure has the attractive property of preserving the linearity of the moment hierarchy, which allows the derivation of analytical results at equilibrium (Bolker and Pacala 1997, 1999). It is quite successful at low densities ($m_1^* \sim 20$) and 1-dimensional systems. However, at intermediate to high densities ($m_1 \sim > 100$) in aggregated patterns, this closure predicts extinction in situations where the point process persists (see dashed-dotted line in Fig. 4a), even for mild correlation regimes. It is nonetheless a useful benchmark result.

The *power-2* closure is obtained as a continuous space analogue to the pair approximation used in discrete spatial systems (Satō and Iwasa 2000),

$$m_3(\xi_1, \xi_2) = \frac{m_2(\xi_1) m_2(\xi_2)}{m_1} + \frac{m_2(\xi_1) m_2(\xi_1 - \xi_2)}{m_1} + \frac{m_2(\xi_1 - \xi_2) m_2(\xi_2)}{m_1} - 2 m_1^3; \quad (22)$$

this closure does predict a persisting population. However, it underestimates quite strongly the second order density, which leads to overshooting the mean density




Fig. 4 Closure comparison. **a** Shows the mean density $\hat{m}_1(t)$ of the point process versus time averaged over 300 sample paths (*rugged black line*). Parameters are $\sigma_B = 0.02$, $\sigma_W = 0.12$, $b = 0.4$, $d = 0.2$, $K = 200$ leading to a segregated pattern of clusters. The *thick dotted black line* shows the predicted mean density from the moment equations with the power-3 or Kirkwood closure, the *dashed black line* corresponds to the power-2 closure and the *dashed-dotted line* corresponds to the power-1 closure. The *dotted grey lines* are the 95 % confidence intervals for the simulation mean value. **b** Shows an estimate the pair correlation function at time $t = 300$ (*thick continuous line*) averaged over 300 independent sample paths, indicating aggregation at short scales, but segregation at intermediate ones. The *dotted thick line* corresponds to the pair correlation function predicted by the solution of the moment hierarchy with the power-3 closure, and the *dashed line* corresponds to the power-2. The 95% confidence intervals for the estimate of the pair correlation function fall within the line

(see Fig. 4b, dashed black line). It is non-linear and thus solutions have to be obtained numerically. There are *asymmetric* versions of this closure that consist of rescaling each additive term in (22) with a set of weighting constants (Law et al. 2003; Murrell et al. 2004). Law and Dieckmann (2000) showed that a particular combination of weighting constants provides a very good fit to simulations. However, this result is difficult to generalize as there is no theory informing how these constants are chosen, they are likely to depend on the details of the model (Murrell et al. 2004; Ovaskainen and Cornell 2006a), and can only be found by comparisons with simulations of the IBM.

Finally, the *power-3* or Kirkwood closure (24) has a distinguished tradition in the statistical mechanics of fluids (Kirkwood and Boggs 1942). Recently, Singer (2004) showed that this closure can be obtained in the hydrodynamic limit after invoking a maximum entropy principle to truncate the BBGKY hierarchy. Earlier motivations for this closure were based on the assumption that each of the pair correlation functions associated with the three edges of the triplet configuration (see Fig. 3) occurs independently of each other *at all spatial scales*,

$$g_3(x_1, x_2, x_3) = g_2(x_1, x_2) g_2(x_1, x_3) g_2(x_2, x_3). \tag{23}$$

Substituting the definition of the k th correlation function in terms of the product densities (17) into (23) for $k = 3$ yields a version of the Kirkwood closure (23) that can be used to close the equation at second order (20)

$$m_3(\xi_1, \xi_2) = \frac{m_2(\xi_1) m_2(\xi_2) m_2(\xi_1 - \xi_2)}{m_1^3}. \quad (24)$$

This closure also underestimates the second order density, but less dramatically so than the power-2 closure, which results in a slightly better prediction of the mean density (see Fig. 4b). Despite its appealing simplicity, the power-3 closure shares the same limitations of the other heuristic closures, e.g. there is no criterion of validity, and it provides poor fit to the equilibrium density even for mildly aggregated patterns (Dieckmann and Law 2000; Raghieb Moreno 2006). Heuristic closures have reasonably good performance in random and segregated spatial configurations, but are significantly more limited in aggregated regimes, with the sole exception of the asymmetric power-2. Their limitation arises from the assumption that there are no irreducible triplet correlations at any scale, in the sense that after fixing a pair that forms an edge, for instance the points x_1 and x_2 (see Fig. 3), the two other edges of the triplet formed with the third point x_3 occur independently of how the first edge is chosen. This can only be true when the three points are sufficiently far apart, but irreducible third order correlations are likely to occur when the three points are close together in aggregated patterns (Fig. 6).

4.2 The Maxent closure

The concept of entropy from an information theoretic point of view, as opposed to the thermodynamical definition of entropy, is tightly related to the uncertainty (or information content) associated with an outcome of a random variable. It can be shown (Khinchin 1957; Shannon and Weaver 1949) that the information content of a particular outcome ($x' + dx'$) of random variable x with probability density $p(x)$, is given by $\log[p(x')dx']$. The entropy functional is constructed by taking the expected value of the information content over all the possible outcomes of x (Jaynes 1957; Khinchin 1957; Shannon and Weaver 1949). To illustrate what this means, consider the uniform distribution on an interval $[a, b] \in \mathbb{R}^+$. It is not surprising that this distribution maximizes the entropy functional if no constraints are introduced, since all the values in its domain of definition have the same probability weight, thus the uncertainty about a specific outcome of a random variable with this distribution is maximal. The opposite situation occurs for the Dirac delta distribution which is centered on one single value, say x' . In this situation, a single value occurs with probability one, and all the others have probability zero, therefore the uncertainty about an outcome of this (pathological) random variable is null.

The principle of maximum entropy is a powerful method that allows the derivation of probability distributions when only but a few average properties are all that is known about a systems. Maximizing the entropy functional subject to the constraints provided by these averages, leads to probability distributions that have the least bias with respect to the known information (Jaynes 1957, 1982; Shannon and Weaver 1949; Khinchin 1957). For instance, maximisation of the entropy constrained to satisfy normalisation and a given mean value leads to the exponential density. Likewise, maximizing the entropy constrained to satisfy normalisation for a given mean and variance leads to

the Gaussian density. For point processes (Daley and Vere-Jones 2004; McFadden 1965) the entropy is defined with respect to some spatial window of observation A , and has two sources of uncertainty, the first is related to the *counts* within A , and the second is related to the *locations* of the n points inside this window. Truncating the hierarchy at order two assumes that only configurations involving up to three points possess irreducible *spatial* information. We carry that assumption forward onto the locational component of the full point process entropy functional, which we then maximise subject to the constraints of normalisation and product densities up to order two, which are given by the truncated hierarchy. We exploit formal relationships between the product densities and the probabilistic objects used to construct the entropy functional of a point process—the *Janossy* densities—that allow the incorporation of the product density constraints onto the entropy functional, and then translate the results of the maximisation procedure in terms of product densities in order to obtain a closure expression.

Our result differs from other maxent closures, like those of Singer (2004) and Hillen (2004), in a number of ways. First, it is *implicit*, in the sense that the *third* order density appears in both sides of the closing expression for truncation at second order. We do so because the Kirkwood closure arises naturally from independence considerations (Singer 2004) for spatial scales larger than the minimum distance for which the pair correlation function is not constant, but it is not valid within the domain of irreducible triplet correlations, i.e. the probability of observing a third point in the triplet depends on how the first two are chosen. If improvements to the Kirkwood closure are to be made, irreducible triplet correlations must appear in the closure. In the maxent method we propose irreducible third order correlations are generated by iteration of the closure relationship, while the first and second order densities, generated by the hierarchy, are held fixed. Second, we assume that these irreducible third order correlations are confined to a finite window, or spatial scale A_0 , which is found by comparison of the normalisation condition for the correlated process with that of a Poisson process of the same mean density. Third, in contrast to other existing approaches, we used all the moments up to the order of the truncation (including the zeroth) to constrain the entropy functional. This is critically important because the zeroth moment is associated with the normalisation constraint, which allows the determination of the domain of triplet correlations.

The variational problem is formulated in terms of the locational entropy functional of the marginal spatial point process. In order to introduce the product densities as constraints, we exploit known expansions of these in terms of the Janossy densities (Daley and Vere-Jones 2003; Janossy 1950) that constitute the probabilistic objects (the likelihoods) required to construct the entropy functional. Whereas Singer (2004) used the k th order product density to constrain an entropy functional, and Hillen (2004), used an L^2 -norm of the moment hierarchy for this purpose, we used instead the classical definition of the entropy functional for a point process, based on the full battery of Janossy densities (Daley and Vere-Jones 2004; McFadden 1965).

The implicit, order two maxent closure (2) resembles the structure of the power-3 or Kirkwood closure (24), but is complemented by a number of correction terms that depend on averages of the product densities over the spatial scale at which triplets are irreducible. Outside this domain, these correction terms vanish and the closure

becomes identical to the power-3. There are two scales of relevance in the closure, one where irreducible triplet correlations are important, and another one where these can be expressed in terms of second and first orders only.

For the sake of completeness, we first discuss known results related to the entropy of spatial point processes in Sect. 4.3, and the key expansions of Janossy densities in terms of product densities. This is followed by the derivation of the implicit maxent closure for truncation at order two (41).

4.3 The entropy of a point process

The Shannon (or information) entropy $H[\mathcal{P}]$ of a stochastic process \mathcal{P} , interpreted as the average uncertainty (or information content) associated with a given outcome of \mathcal{P} , is defined as minus the expected value of the log-likelihood L (Daley and Vere-Jones 2003, 2004; Jaynes 1957, 1982; Khinchin 1957; Shannon and Weaver 1949),

$$H[\mathcal{P}] = -E\{\log(L)\}. \quad (25)$$

The specialisation of the entropy (25) to point processes requires a special form of the likelihood, given that in a realisation of a point process of the form $\{x_1, \dots, x_n\}$ in a window A there are two sources of uncertainty. The first comes from uncertainty about the number of points n within A (the counts), which is controlled by an integer-valued probability distribution $p_n = \Pr\{N(A) = n\}$. Conditionally on the value of n , the other contribution comes from the uncertainty associated with the *locations* of the n points, which is given by a symmetric (in the sense of invariance under permutations of the indices) probability density $s_n(x_1, \dots, x_n|A)$ on $A^{(n)}$. Thus, the likelihood of a spatial point process is the probability of finding n points within A , each in one of the infinitesimal locations dx_1, \dots, dx_n and nowhere else within A . This coincides with the definition of the local Janossy density (Daley and Vere-Jones 2003, 2004; Janossy 1950)

$$L_A(x_1, \dots, x_n) = p_n s_n(x_1, \dots, x_n|A) = j_n(x_1, \dots, x_n|A). \quad (26)$$

Separating the contributions due to the counts and those due to spatial information, we can represent the entropy of a point process \mathcal{N}_A on a window A by Daley and Vere-Jones (2003, 2004)

$$H[\mathcal{N}_A] = - \sum_{r=0}^{\infty} p_r \log(r! p_r) - \sum_{r=1}^{\infty} p_r \int_{A^{(r)}} s_r(x_1, \dots, x_r) \log[s_r(x_1, \dots, x_r)] dx_1 \dots dx_r, \quad (27)$$

where the integrals calculate the contribution due to the locations, and the sums that of the counts. If we fix the expected number of points in A , $\mu = m_1|A| = E[N(A)]$, it can be shown that the first sum in (27) is maximized by the Poisson distribution (Daley and Vere-Jones 2004; Khinchin 1957; McFadden 1965),

$$p_r = \frac{\mu^r}{r!} \exp(-\mu).$$

Conditional on the counts r , the second sum is maximized by the uniform density on $A^{(r)}$

$$s_r \equiv \frac{1}{|A|^r}.$$

Thus, the point process of maximum entropy is the homogeneous Poisson point process with first order density m_1 (Daley and Vere-Jones 2008, 2004). For closure purposes we use the definition (25) written in terms of the local Janossy densities

$$H[\mathcal{N}_A] = - \sum_{n=0}^{\infty} \frac{1}{n!} \int_{A^{(n)}} j_n(x_1, \dots, x_n | A) \log[j_n(x_1, \dots, x_n | A)] dx_1 \dots dx_n, \tag{28}$$

where division by $n!$ ensures normalisation with respect to the $n!$ permutations of the n indices. Our method of closure consists of maximizing (28) constrained to satisfy the product densities up to the order of truncation. These can only be meaningfully incorporated as constraints if they can be expressed in terms of integrals over A of the Janossy densities. We do this by using the expansion (Daley and Vere-Jones 2003),

$$m_n(x_1, \dots, x_n) = \sum_{q=0}^{\infty} \frac{1}{q!} \int_{A^{(q)}} j_{q+n}(x_1, \dots, x_q, y_1, \dots, y_n) dy_1 \dots dy_n, \tag{29}$$

where the inverse relationship,

$$j_n(x_1, \dots, x_n | A) = \sum_{q=0}^{\infty} \frac{(-1)^q}{q!} \int_{A^{(q)}} m_{n+q}(x_1, \dots, x_n, y_1, \dots, y_q) dy_1 \dots dy_q, \tag{30}$$

can be used to translate the results of the constrained optimisation procedure in terms of product densities in order to yield a closure for the product density hierarchy.

4.4 Maximum entropy closure at order $k = 2$

In the case of the non-homogeneous Poisson point process, which maximizes the entropy functional (28), all the points can in principle depend on the specific locations, but these are uncorrelated. For this special case the expansion of the likelihoods in terms of the product densities (30) takes the simplified form,

$$j_n(x_1, \dots, x_n | A) = \prod_{p=1}^n m_1(x_p) \sum_{q=0}^{\infty} \frac{(-1)^q}{q!} \prod_{l=0}^q m_1(y_l) |A|^l. \tag{31}$$




Fig. 5 Estimated radial pair correlation functions at equilibrium $g_2^*(r)$ from simulations of the point process in Sect. 2.3 with dispersal and mortality kernels given by symmetric bivariate Gaussians. Parameters ($\sigma_B = 0.04, \sigma_W = 0.04$) lead to a mildly aggregated pattern (case b *dashed-dotted line*) and a segregated pattern of clusters [case a *continuous line*, ($\sigma_B = 0.02, \sigma_W = 0.12$)]. In **b** we note that correlations decay quickly and become constant at a spatial lag $r > 0.2$, whereas in **a** there are distinct patterns in at least two spatial scales. Aggregation in the smaller ones, and segregation at intermediate ones. Case (c) (*dotted line*) corresponds to a Poisson point process

If the process is a spatially stationary and homogeneous Poisson point process, then all the product densities become simple powers of the mean density (Diggle 1983; Daley and Vere-Jones 2003), which further simplifies (30) to,

$$j_n(x_1, \dots, x_n | A) = m_1^n \exp(-m_1 |A|). \tag{32}$$

From (30) we have that the probability of observing n points within a window A is

$$\Pr [N(A) = n] = \frac{1}{n!} \int_{A^{(n)}} j_n(x_1, \dots, x_n | A) dx_1 \cdots dx_n, \tag{33}$$

which after substituting (32) into (33) leads to the Poisson distribution

$$\Pr [N(A) = n] = \frac{(m_1 |A|)^n \exp(-m_1 |A|)}{n!}.$$

We assume somewhat crudely that the Janossy expansions of the point process associated with the moment hierarchy have an intermediate structure between the two extreme cases (30) where the spatial configurations of all orders are irreducible, and the Poisson point process (32) where all the locations occur independently. This can be justified from the truncation assumption, since truncating the hierarchy at order two implicitly assumes that terms of order equal or higher than four do not contribute to the formation of second and third order spatial correlations. Also we see in estimates of the pair correlation functions for the point process discussed in Sect. 2, shown in Fig. 5 that there is a region in the parameters for which the spatial correlations of second order decay quickly. Case (a) corresponds to segregated clusters and thus the pair correlation oscillates around one. There are two different scales with pattern there,

the first is associated with the clusters (the region where $g_2 > 1$) and another with the separation between the clusters themselves ($g_2 < 1$). Case (b) on the other hand corresponds to a simply aggregated pattern. In this latter case we see clearly that there is a spatial scale for which the pair correlation function becomes constant and identical to one, therefore

$$m_2(r) = m_1^2, \quad r \gg r_0$$

for some spatial scale r_0 . This assumption is tantamount to requiring that the Janossy expansions of the process to have the form,

$$\begin{aligned}
 j_n(x_1, \dots, x_n | A) = & \sum_{q=0}^{k+1-n} \frac{(-1)^q}{q!} \int_{A^{(n)}} m_{n+q}(x_1, \dots, x_n, y_1, \dots, y_q) dy_1 \dots dy_q \\
 & + \prod_{p=1}^n m_1(x_p) \sum_{q>k+1-n}^{\infty} \frac{(-1)^q}{q!} \int_{A^{(q)}} \prod_{r=1}^q m_1(y_r) dy_r, \quad (34)
 \end{aligned}$$

where the first term corresponds to the terms that make contributions due to spatial correlations, and the second term is the (non-homogeneous) Poisson remainder. The closure assumption implies that only the Janossy densities of order up to $k + 1$ make contributions to the locational entropy, in which case the entropy functional (28) becomes

$$\begin{aligned}
 H_{loc}^{(3)}[\mathcal{N}_A] = & -J_0(A) \log[J_0(A)] - \sum_{n=1}^3 \sum_{1 \leq i_1 < \dots < i_n \leq 3} \frac{(3-n)!}{3!} \\
 & \times \int_{A^{(n)}} j_n(x_{i_1}, \dots, x_{i_n} | A) \log[j_n(x_{i_1}, \dots, x_{i_n} | A)] dx_{i_1} \dots dx_{i_n} \quad (35)
 \end{aligned}$$

where $J_0(A)$ is the avoidance probability in A . The first constraint added to (36) is that of normalisation,

$$1 = \sum_{n=0}^{\infty} \frac{1}{n!} \int_{A^{(n)}} j_n(x_1, \dots, x_n) dx_1 \dots dx_n,$$

which after simplification with the assumption (34) can be added to the entropy functional

$$\begin{aligned}
 + \Lambda_0 \cdot & \left(J_0(A) + \sum_{q=1}^3 \sum_{1 \leq i_1 < \dots < i_q \leq 3} \frac{(3-q)!}{3!} \int_{A^{(q)}} j_q(x_{i_1}, \dots, x_{i_q} | A) dx_{i_1} \dots dx_{i_q} \right. \\
 & \left. + \sum_{n>3}^{\infty} \prod_{i=1}^n m_1(x_i) \sum_{l>3-n}^{\infty} \frac{(-1)^l}{l!} \prod_{r=1}^l \int_{A^{(r)}} m_1(y_r) dy_r - 1 \right)
 \end{aligned}$$

where Λ_0 is a (constant) Lagrange multiplier. The second constraint is that of the first order product density $m_1(x_i)$

$$\begin{aligned}
 &+ \sum_{1 \leq i_1 \leq 3} \frac{1}{3} \int_A \Lambda_1(x_{i_1}) \left(\sum_{q=0}^2 \sum_{1 \leq i_1 < \dots < i_q \leq 3} \frac{(3-q)!}{3!} \right. \\
 &\quad \left. \times \int_{A^{(q)}} j_{1+q}(x_{i_1}, \dots, x_{i_n}, y_{i_1}, \dots, y_{i_q} | A) dy_{i_1} \dots dy_{i_q} - m_1(x_{i_1}) \right) dx_{i_1}. \tag{36}
 \end{aligned}$$

where $\Lambda_1(x_{i_1})$ is a vector of functional Lagrange multipliers, each associated with the permutations in the locations x_1, x_2 and x_3 comprising the triplet. Finally, the constraint for the second order product density $m_2(x_{i_1}, x_{i_2})$ is

$$\begin{aligned}
 &+ \sum_{1 \leq i_1 < i_2 \leq 3} \frac{1}{6!} \int_{A^{(2)}} \Lambda_2(x_{i_1}, x_{i_2}) \left(\sum_{q=0}^1 \sum_{1 \leq i_1 < \dots < i_q \leq 3} \frac{(3-q)!}{3!} \right. \\
 &\quad \left. \times \int_{A^{(q)}} j_{2+q}(x_{i_1}, \dots, x_{i_n}, y_{i_1}, \dots, y_{i_q} | A) dy_{i_1} \dots dy_{i_q} - m_2(x_{i_1}, x_{i_2}) \right) dx_{i_1} dx_{i_2}. \tag{37}
 \end{aligned}$$

Likewise, the $\Lambda_2(x_{i_1}, x_{i_2})$ are the Lagrange multipliers associated with each of the permutations of the pairs in the triplet. The Euler–Lagrange equations of the functional (35)–(37) are

$$\begin{aligned}
 \frac{\delta H^{(3)}}{\delta J_0(A)} &= -1 - \log[J_0(A)] + \Lambda_0 = 0, \\
 \frac{\delta H^{(3)}}{\delta j_1(x_{i_1})} &= -\frac{1}{3} (1 + \log j_1 [(x_{i_1})]) + \frac{1}{3} \Lambda_0 + \frac{1}{3} \Lambda_1(x_{i_1}) = 0, \quad 1 \leq i_1 \leq 3 \\
 \frac{\delta H^{(3)}}{\delta j_2(x_{i_1}, x_{i_2})} &= -\frac{1}{6} (1 + \log [j_2(x_{i_1}, x_{i_2})]) + \frac{1}{6} \Lambda_0 + \frac{1}{3} \Lambda_1(x_{i_1}) \\
 &\quad + \frac{1}{6} \Lambda_2(x_{i_1}, x_{i_2}) = 0, \quad 1 \leq i_1 \leq i_2 \leq 3 \tag{38} \\
 \frac{\delta H^{(3)}}{\delta j_3(x_1, x_2, x_3)} &= -\frac{1}{6} (1 + \log [j_3(x_1, x_2, x_3)]) + \frac{1}{6} \Lambda_0 + \frac{1}{2} [\Lambda_1(x_1) + \Lambda_1(x_2) \\
 &\quad + \Lambda_1(x_3)] + \frac{1}{2} [\Lambda_2(x_1, x_2) + \Lambda_2(x_2, x_3) + \Lambda_2(x_1, x_3)] = 0.
 \end{aligned}$$

It can be seen by inspection that each of the second variations is inversely proportional to minus the Janossy density of order k . Since these are all probability densities, each of the second variations is negative and thus the extrema given in the first variation (39)

are maxima. Solving the Euler–Lagrange equations (39) for the Lagrange multipliers yields

$$\begin{aligned}
 \Lambda_0 &= 1 + \log[J_0(A)] \\
 \Lambda_1(x_1) &= \log \left[\frac{j_1(x_1)}{J_0(A)} \right] \\
 \Lambda_1(x_2) &= \log \left[\frac{j_1(x_2)}{J_0(A)} \right] \\
 \Lambda_1(x_3) &= \log \left[\frac{j_1(x_3)}{J_0(A)} \right] \\
 \Lambda_2(x_1, x_2) &= \log \left[\frac{J_0(A) j_2(x_1, x_2)}{j_1^2(x_1)} \right] \\
 \Lambda_2(x_2, x_3) &= \log \left[\frac{J_0(A) j_2(x_1, x_3)}{j_1^2(x_2)} \right] \\
 \Lambda_2(x_1, x_3) &= \log \left[\frac{J_0(A) j_2(x_2, x_3)}{j_1^2(x_3)} \right].
 \end{aligned}
 \tag{39}$$

Substituting the Lagrange multipliers in (39) into the equation for the first variation with respect to j_3 in (39) yields an expression that relates the Janossy density of third order to the lower order ones under the assumption of maximum entropy constrained by the moments, namely

$$j_3(x_1, x_2, x_3|A) = \frac{j_2(x_1, x_2|A) j_2(x_2, x_3|A) j_2(x_1, x_3|A)}{j_1(x_1|A) j_1(x_2|A) j_1(x_3|A)} J_0(A),
 \tag{40}$$

Equation (40) is formally similar to the Kirkwood closure. However, there are a number of important differences. First, it varies with the choice of the window A , since it depends on the *local* likelihoods (see Fig. 6) rather than the product densities used in the Kirkwood closure, which are global properties that do not depend on the window of observation. This domain A depends on the spatial scale for which the third particle in the triplet becomes independent of the other two. Second, the closure is weighted by the avoidance probability $J_0(A)$. This term is conceptually similar to the exponential weight suggested by Meeron (1957) and Salpeter (1958), but now arises from a maximum entropy consideration. The relationship (40) can be used as a closure of the moment hierarchy after using the expansions (30) and (34) that allow the Janossy densities to be expressed in terms of product densities.

Since the underlying point process is spatially stationary by construction, then the mean density is constant, and the densities of higher orders depend on the relative rather than absolute distances between points. After rescaling the product densities in the expansion by the area of the window A (the product densities that come from the hierarchy are defined in terms of the much larger spatial window used to observe the




Fig. 6 The domain A represents the region beyond which a third particle becomes independent of the other two. Shifting x'_3 to x_3 , makes that third point independent of the other two. This corresponds to the spatial scale for which the assumptions leading to the Kirkwood closure are valid

full process) we have that the maxent closure is given by if $|\xi_1| \leq r_0$ and $|\xi_2| \leq r_0$ and $|\xi_2 - \xi_1| \leq r_0$

$$\begin{aligned}
 m_3(\xi_1, \xi_2) = & \left[m_2(\xi_1) - |A_0| \int_{A_0} m_3(\xi_1, \xi'_2) d\xi'_2 \right] \\
 & \times \left[m_2(\xi_2) - |A_0| \int_{A_0} m_3(\xi_2, \xi_2 - \xi'_1) d\xi'_1 \right] \\
 & \times \left[m_2(\xi_2 - \xi_1) - |A_0| \int_{A_0} m_3(\xi_2 - \xi'_1, \xi'_1) d\xi'_1 \right] \\
 & \times \frac{J_0(A_0)}{\left[m_1 - |A_0| \int_{A_0} m_2(\xi'_1) d\xi'_1 + \frac{|A_0|^2}{2} \int_{A_0^{(2)}} m_3(\xi'_1, \xi'_2) d\xi'_1 d\xi'_2 \right]^3}, \tag{41}
 \end{aligned}$$

else

$$m_3(\xi_1, \xi_2) = \frac{m_2(\xi_1) m_2(\xi_2) m_2(\xi_2 - \xi_1)}{m_1^3} \tag{42}$$

where the circular domain A_0 of radius r_0 is determined from the normalisation constraint (described below). The avoidance function $J_0(A_0)$ is given by

$$\begin{aligned}
 J_0(A_0) = & 1 - m_1|A_0| + \frac{|A_0|}{2} \int_{A_0} m_2(\xi_1)d\xi_1 - \frac{|A_0|}{6} \int_{A_0^{(2)}} m_3(\xi_1, \xi_2)d\xi_1d\xi_2 \\
 & + \sum_{n=4}^{\infty} \frac{(-1)^n}{n!} (m_1|A_0|)^n
 \end{aligned}
 \tag{43}$$

and the summation term is equal to

$$\sum_{n=4}^{\infty} \frac{(-1)^n}{n!} (m_1|A_0|)^n = \exp(-m_1|A_0|) - 1 + m_1|A_0| - \frac{(m_1|A_0|)^2}{2} + \frac{(m_1|A_0|)^3}{6}.$$

After simplifying we have

$$\begin{aligned}
 J_0(A_0) = & \exp(-m_1|A_0|) + \frac{|A_0|}{2} \int_{A_0} m_2(\xi_1)d\xi_1 - \frac{(m_1|A_0|)^2}{2} \\
 & - \frac{|A_0|}{6} \int_{A_0^{(2)}} m_3(\xi_1, \xi_2)d\xi_1d\xi_2 + \frac{(m_1|A_0|)^3}{6}.
 \end{aligned}
 \tag{44}$$

In order to obtain the set A_0 in the correction terms of the closure, we first need to identify the spatial scale r_0 beyond which two points become independent. This is equivalent to finding the smallest region A_0 for which the correlated point process has the same statistics of a Poisson process of the same mean density. This domain is obtained by comparing the avoidance functions for each case, which must coincide for this specific set. Since the avoidance probability for a homogeneous Poisson point process of intensity m_1 for some reference window B is equal (Daley and Vere-Jones 2003) to

$$J_0^*(B) = \exp(-m_1|B|),
 \tag{45}$$

Thus the set A_0 must satisfy

$$J_0(A_0) = J_0^*(A_0).
 \tag{46}$$

Substituting the RHS of (45) and (44) into (46) leads to the integral equation

$$\int_{A_r} m_2(\xi_1)d\xi_1 - m_1^2|A_r| - \frac{1}{3} \int_{A_r^{(2)}} m_3(\xi_1, \xi_2)d\xi_1d\xi_2 + \frac{m_1^3|A_r|^2}{3} = 0,
 \tag{47}$$

where $A_r = B(0, r)$ is the ball of radius r centered at the origin. Since all the product densities are given by the hierarchy and the closure relationship (41), the only unknown in (47) is the domain A_0 that satisfies the equality (47). This can be found by evaluating the RHS of (47) for an increasing family of domains A_r . The values of r that satisfy the equality are the roots of interest. There are four possible scenarios for these roots:

1. The trivial root, $r = 0$ is the only solution. This is always a solution by simple inspection.
2. A single non-trivial root r^* .
3. A finite number of n non trivial roots $r_1^*, r_2^*, \dots, r_n^*$.
4. An infinite number of roots.

A criterion of validity for the closure scheme can be built on the basis of the number of roots. Case 1 indicates that there is not a scale within the observed range of r for which correlations decay as powers of the mean density, and thus truncation should be tried at a higher order. Case 2 indicates that there is a single Poisson domain A_0 and thus the closure assumptions are consistent with the predicted values of the hierarchy. Case 3 indicates that there are several scales of spatial pattern, due to correlations that oscillate as they decay, i.e. segregated clusters (see Fig. 5). In this situation each scale of pattern should be treated separately. An infinite number of roots (case 4) indicates that the process is indistinguishable from a Poisson process at all scales.

Although the closure expression seems complicated, we note that the integral correction terms in the numerator and denominator are of similar magnitude, and relatively small in comparison with the correction introduced by avoidance probability, which by far dominates the closure. In this situation we have a much simpler approximation to the exact closure, given by

$$m_3(\xi_1, \xi_2) \approx \frac{m_2(\xi_1) m_2(\xi_2) m_2(\xi_2 - \xi_1)}{m_1^3} \exp(-m_1 |A_0|). \quad (48)$$

5 Numerical implementation

The numerical solution of the hierarchy with the maxent closure requires two separate modules of code: one for the integration of the hierarchy itself, and the other for the iterative procedure that computes the third order density. The first, which we call the ‘outer’ code, consists of a standard numerical integration scheme that predicts the first and second order product densities at a time $(t + h)$ using the first, second and third order ones at time t as input, where h is a small time step. The second module, or ‘inner code’, computes the third order density at time $(t + h)$ from the maxent closure. The inner code starts by computing an initial value for the area of normalisation $A_0^{(old)}$ using the values of the first and second order densities at time $(t + h)$, and the third order density at time t as an initial trial. This first value $A_0^{(old)}$ is then substituted in the maxent closure expression (41) to produce an updated value for the third order density. The area of normalisation is recalculated with the updated third order density to produce a new value $A_0^{(new)}$; if the relative difference between the old and the new

radii associated with each normalisation area falls below some pre-specified tolerance, then the iteration stops and the final value of the third order density at time $(t + h)$ is the one being used to calculate the last iteration of area of normalisation. If not, the iterations continue until the tolerance is achieved. We now propose an algorithm for the implementation the maxent closure, and subsequently show its performance for a broad range of parameters of the spatial scales. Our numerical results are well behaved and convergence of the iteration scheme occurs rapidly for a sufficiently small time step ($h = 0.1$), where typically two or three iterations of the closure are sufficient for a relative error tolerance within one percent. The problem consists of solving the coupled system

$$\left\{ \begin{aligned} \frac{d}{dt} m_1(t) &= r m_1(t) - d_N \int_B W(\xi_1) m_2(\xi_1, t) d\xi_1 \\ \frac{1}{2} \frac{d}{dt} m_2(\xi_1, t) &= b \int_B B(\xi_2) m_2(\xi_1 - \xi_2, t) d\xi_2 + b B(\xi_1) m_1(t) - d m_2(\xi_1, t) \\ &\quad - d_N W(\xi_1) m_2(\xi_1, t) - d_N \int_B W(\xi_2) m_3(\xi_1, \xi_2, t) d\xi_2. \end{aligned} \right. \tag{49}$$

where

$$B \subset \mathbb{R}^2, \quad t \in [0, T_{max}] \subset \mathbb{R}_0^+,$$

with initial condition

$$m_1(0) = n_0, \quad m_2(\xi_1, 0) = n_0^2, \quad m_3(\xi_1, \xi_2, 0) = n_0^3.$$

The window B should be large enough to approximate correctly the integral terms, which are originally defined over all the plane. This hierarchy can be closed at order 2 with the maxent closure (41)

$$\begin{aligned} m_3(\xi_1, \xi_2) &= \left[m_2(\xi_1) - |A_0| \int_{A_0} m_3(\xi_1, \xi'_2) d\xi'_2 \right] \\ &\times \left[m_2(\xi_2) - |A_0| \int_{A_0} m_3(\xi_2, \xi'_1) d\xi'_1 \right] \\ &\times \left[m_2(\xi_2 - \xi_1) - |A_0| \int_{A_0} m_3(\xi_2 - \xi'_1, \xi'_1) d\xi'_1 \right] \\ &\times \frac{J_0(A_0)}{\left[m_1 - |A_0| \int_{A_0} m_2(\xi'_1) d\xi'_1 + \frac{|A_0|^2}{2} \int_{A_0 \times A_0} m_3(\xi'_1, \xi'_2) d\xi'_1 d\xi'_2 \right]^3}, \end{aligned} \tag{50}$$

which is applied if each the three distance vectors $(\xi_1, \xi_2$ and $\xi_2 - \xi_1$, see Fig. 6) connecting the three points in the triple configuration fall within the normalisation domain A_0 . Outside of this region we apply the Kirkwood closure on the basis of probabilistic independence of the third point in the triplet, as discussed in the previous section

$$m_3(\xi_1, \xi_2) = \frac{m_2(\xi_1) m_2(\xi_2) m_2(\xi_2 - \xi_1)}{m_1^3}. \tag{51}$$

In the maxent closure (50) the avoidance function $J_0(A_0)$ is given by

$$J_0(A_0) = \exp(-m_1|A_0|).$$

The circular domain A_0 is computed from the comparison between the normalisation constraint for the truncated hierarchy and that of a Poisson process of the same mean intensity. It is calculated by finding the value of r that satisfies

$$\int_{A_r} m_2(\xi'_1) d\xi'_1 - m_1^2|A_r| - \frac{1}{3} \int_{A_r^{(2)}} m_3(\xi'_1, \xi'_2) d\xi'_1 d\xi'_2 + \frac{m_1^3|A_r|^2}{3} = 0. \tag{52}$$

where A_r is the 2-dimensional ball of radius r centred at the origin.

5.1 Algorithm for the numerical implementation

The coupled system of product density equations with the maxent closure can be solved from the following algorithm:

1. From a sequence of radii $r_i = 0, \dots, r_{max}$, construct an increasing family of domains A_{r_i} .
2. At time $t = 0$ the initial configuration is given by a homogeneous Poisson point process, thus all the product densities are powers of the mean density $N_0/|X|$, where X is the computational spatial arena, and N_0 is the population size at time $t = 0$.
3. While the elapsed time $t < T_{max}$ do
 - (a) Integrate forward the densities $m_1(t+h)$ and $m_2(\xi_1, t+h)$ from the hierarchy using a standard numerical procedure.
 - (b) Use the value of the triplet density at the earlier time step $m_3^{(old)}(\xi_1, \xi_2, t)$ as the initial guess in the normalisation condition for the Poisson area A_0 . Generate a sequence of values $f(r_i)$ by calculating the normalisation condition (52) for each the domains previously constructed in Step 1 according to

$$f(r_i) = \int_{A_{r_i}} m_2(\xi'_1, t+h) d\xi'_1 - \frac{1}{3} \int_{A_{r_i}^{(2)}} m_3^{(old)}(\xi'_1, \xi'_2, t) d\xi'_1 d\xi'_2 - m_1^2(t+h) a_{r_i} + \frac{1}{3} m_1^3(t+h) a_{r_i}^2 \tag{53}$$

where the a_{r_i} are the areas for each of the A_{r_i} .

- (c) Find the largest value r_o that satisfies $f(r_o) = 0$ by linear interpolation between the consecutive r_i where $f(r_i)$ changes sign.
- (d) Use r_o from Step 3c to generate the estimate of the Poisson domain $A_0 = A_{r_o}$.
- (e) Loop the spatial arguments ξ_1 and ξ_2 over the computational spatial arena.
- (f) Compute the magnitudes d_1, d_2 and d_3 of the distance vectors ξ_1, ξ_2 and $\xi_2 - \xi_1$
- (g) if $d_1 \leq r_o$ and $d_2 \leq r_o$ and $d_3 \leq r_o$ apply the maxent closure

$$m_3^{(new)}(\xi_1, \xi_2) = \frac{\exp(-m_1 |A_0|)}{\left[m_1 - A_0 \int_{A_0} m_2(\xi'_1) d\xi'_1 + \frac{A_0^2}{2} \int_{A_0}^{(2)} m_3^{(old)}(\xi'_1, \xi'_2) d\xi'_1 d\xi'_2 \right]^3} \times \left[m_2(\xi_1) - A_0 \int_{A_0} m_3^{(old)}(\xi'_1, \xi'_2) d\xi'_2 \right] \times \left[m_2(\xi_2) - A_0 \int_{A_0} m_3^{(old)}(\xi_2, \xi_2 - \xi'_1) d\xi'_1 \right] \times \left[m_2(\xi_2 - \xi_1) - A_0 \int_{A_0} m_3^{(old)}(\xi_2 - \xi'_1, \xi'_1) d\xi'_1 \right],$$

- (h) else use the Kirkwood closure

$$m_3^{(new)}(\xi_1, \xi_2) = \frac{m_2(\xi_1) m_2(\xi_2) m_2(\xi_2 - \xi_2)}{m_1^3}. \tag{54}$$

- (i) Recompute the Poisson domain $A_0^{(new)}$ and its radius $r_o^{(new)}$ by inserting the corrected triplet density $m_3^{(new)}$ from Step 3e into the normalisation equation into Step 3c and estimate a new root r_n .
- (j) If the difference between the old radius and the new one falls within the error tolerance

$$\frac{|r_o - r_o^{(new)}|}{r_o} \leq \text{tolerance}$$

then the third order density at time $(t + h)$ is the one calculated at Step 3e

$$m_3(\xi_1, \xi_1, t + h) = m_3^{(new)}(\xi_1), \xi_2$$

else the old third order density becomes the new third order density

$$m_3^{(new)} \rightarrow m_3^{(old)}$$

and repeat Steps 3c through 3i until the error falls within the tolerance.




Fig. 7 Comparison between the mean density (*thick line*) for a sample of 300 simulations of the point process for the mildly aggregated case $\sigma_B = 0.05$, $\sigma_W = 0.05$ and the truncated product density hierarchy using various closures. The the maximum entropy closure (maxent) (*continuous black line*), the power-3 (*dashed*), the symmetric power-2 (*dotted*) and power-1 (*dashed-dotted*). The maximum entropy closure provides the best fit to the equilibrium values of the IBM. However the performance of all the closures is poor during the transient regime

4. update the elapsed time

$$t \rightarrow t + h.$$

5.2 Closure performance

We applied the simulation algorithm¹ introduced in the previous Sect. 5.1 using a spatial discretisation of 47 points per linear dimension, and the domain B was the unit square $[-1/2, 1/2] \times [-1/2, 1/2]$. The spatial integrals were computed using the trapezoidal rule, and the convolution in (49) was calculated using the fast Fourier transform. For the solution of the moment hierarchy we use a fourth-order Runge–Kutta scheme (with a time step $h = 0.1$). Convergence was checked by halving the time step and the spatial discretisation and no significant differences were found ($m_1^* = 168.6$, $\Delta x = 1/47$, $h = 0.1$ and $m_1^* = 168.9$, $\Delta x = 1/95$, $h = 0.05$, for $\sigma_W = \sigma_B = 0.05$).

The maxent closure is expected to work well in situations where the spatial scales of dispersal and mortality are similar, since this combination of parameters tends to produce a single scale of spatial pattern of mild aggregation (see Fig. 5), where higher order terms are small. Figure 7 compares the dynamics of the mean density predicted

¹ Code available from first author on request.

by the maxent closure in a mildly aggregated regime ($\sigma_B = \sigma_W = 0.05$) against averages of the point process model and the other closure methods used in the literature, power-1, power-2 and power-3 (but the *asymmetric* power-2 is not used in the comparison). We see that the maxent closure outperforms the other closures. As before, in all cases the transient is predicted poorly. This is to be expected of the maxent method, because the locational entropy can be assumed to be maximised only once the stochastic process has reached its stationary distribution. For this reason, even with the correction terms, the truncated hierarchy with the maxent closure fails at capturing the transient behavior, which typically consists of long range spatial correlations that decay only once the density-dependent mortality term is large enough to cause mixing at longer scales, thus producing a shorter correlation scale.

The ability of the maxent closure to predict accurately the mean density changes dramatically when the two interaction kernels have very different characteristic scales. This combination of parameters leads to several scales of pattern, that can consist of short range aggregation compensated by long range segregation, or short scale segregation compensated by long range clustering. This occurs because the total number of pairs over sufficiently long ranges must be equal to the density squared. Thus, extreme aggregation over short scales must be compensated by segregation over the longer scales in order to preserve the total number of pairs. When dispersal has a much shorter characteristic scale than that of density-dependent mortality, the resulting pattern consists of segregated clusters. This situation violates the closure assumptions (that require a single scale of pattern), and we expect the validity checks in the maxent closure to be activated in this situation. This is illustrated for two types of aggregated patterns in Fig. 8. The upper three panels correspond to segregated clusters ($\sigma_B = 0.02$, $\sigma_W = 0.12$), and the lower three to the mild aggregation case discussed earlier ($\sigma_B = \sigma_W = 0.04$). The left column conformed by Figs. 8a, b show typical point patterns obtained at the same time at which the numerical solution of the hierarchy stopped, $t = 1.56$ in Fig. 8a, because of the validity check, and $t = 80$ in Fig. 8b which was long enough to reach equilibrium. The center column, consisting of Fig. 8c, d, displays kernel density estimates of the pair correlation function for the point patterns shown to the left. We see in Fig. 8c a very high degree of aggregation at short scales followed by long range segregation. Finally, Fig. 8e, f shows the dynamics of the area of correlations $A_0(t)$ for both regimes. We see failure of the maxent closure to find a non-trivial root for A_0 in Fig. 8e after a short transient, as should be expected due to the presence of various scales of pattern detected in the pair correlation function in Fig. 8c. In this situation, the extreme form of ‘checkerboard’ aggregation requires truncation at a higher order. Since the pair correlation function is clearly not constant, but yet the normalisation constraint only finds the trivial root zero, the validity check is activated and the numerical solution of the hierarchy stops. By contrast, in the lower panels when the degree of clustering is comparatively smaller, the method succeeds in finding a single root A_0 that eventually reaches a single equilibrium (see Fig. 8f).

We carried out a systematic exploration of the behavior of the maxent closure for a wide range of combinations (441 in total) of the spatial parameters falling within the range $[0.02, 0.12]$ that correspond to those explored earlier by Law et al. (2003), and compare the results with the predictions of the point process, and the product density hierarchy with the power-3 closure. This allows the assessment of the




Fig. 8 Behavior of the area of corrections in the maxent closure for two types of aggregated spatial patterns. The upper three panels correspond to a segregated pattern of clusters with $\sigma_B = 0.02$, $\sigma_W = 0.12$, and the lower panels to a mildly aggregated pattern with $\sigma_B = \sigma_W = 0.04$. The left column shows a single point pattern at the end of the simulation, the middle column shows a kernel density estimate of the pair correlation function for the pattern displayed in the left and the right column shows the temporal behavior of the area of the set in the correction terms




Fig. 9 Simulation results of the product density hierarchy with the maxent for various values of the characteristic spatial scales of dispersal σ_B (horizontal axis) and mortality σ_W (vertical axis). **a** Shows the equilibrium mean density m_1^* . **b** Shows the value of the second order product density at equilibrium evaluated at the origin, normalized by the squared mean density. In this panel values higher than one indicate clustering at short scales, and values below one indicate segregation. **c** Shows the value at equilibrium of the area of the domain used in the correction terms A_0

relative importance of the correction terms in the maxent closure. The upper limit in the parameter domain was chosen because for that scale ($\sigma_B = \sigma_W = 0.12$) there is only a very small departure from complete spatial randomness. Figure 9 shows various equilibrium values predicted by the product density hierarchy with the maxent closure. Figure 9a corresponds to the mean density, Fig. 9b shows the equilibrium value of the second moment at the origin, normalized by the mean density squared, and finally, Fig. 9c shows the area of normalisation at equilibrium. The removed regions (white) in Fig. 9a result from the application of the validity check of normalisation, since for this parameter the area of correlations is zero (see Fig. 9c), but the second order product density indicates the existence of spatial pattern.




Fig. 10 Comparison of the mean density m_1^* at equilibrium predicted by an ensemble average of the point process model (a), the maxent closure (b), and the power-3 or Kirkwood closure (c). In b, the white region in the upper left corner corresponds to the domain where the normalisation constraint returns a trivial root for values of the second order product density that indicate the presence of spatial pattern, activating the validity check (47)

In Fig. 10 we compare the mean equilibrium density predicted from an average of the space–time point process (a), the maxent closure (b), and the power-3 closure (c). The maxent closure is not a good predictor of the mean density for intermediate to low ranges of mortality combined with long range dispersal; in this regime both the qualitative and quantitative behavior of the closure is poor. We see a sharp drop in the values of the mean density, whereas in the point process model it grows monotonically before reaching the plateau that occurs when both dispersal and mortality act over long scales. This combination of parameters leads to segregation at short scales and long range (albeit mild) aggregation. The maxent method detects only the scale of aggregation, which produces comparatively larger values of the area of correlations (see Fig. 9c). This leads to over-correction in the maxent closure, which results in an equilibrium density that falls well below that predicted by the point process model. In this regime, the power-3 closure provides a much more precise prediction of the equilibrium density, both qualitatively and quantitatively. For sufficiently short ranges of dispersal together with short to intermediate ranges of mortality the point process model predicts extinction, as already noted earlier by Law and Dieckmann (2000) and Law et al. (2003). In this regime, neither the maxent closure nor the power-3 closure is capable of predicting the persistence/extinction threshold, and the maxent validity check does not seem to operate either. However, for intermediate ranges of aggregation close or above the main diagonal ($\sigma_W = \sigma_B$), the maxent closure does provide an improved prediction of the equilibrium density, with the added benefit of the criterion of validity being activated when dispersal is short range with long range mortality, which leads to different scales of pattern (Fig. 11).

We computed the relative error between the equilibrium density of the point process, and that predicted by the moment equations with the two closures, shown in Fig. 11. Figure 11a corresponds to the maxent closure and Fig. 11b to the power-3. We see that the maxent closure has larger relative error than the power-3 for values located below the diagonal ($\sigma_W = \sigma_B$), which are associated with segregated spatial patterns (see Fig. 9b). In contrast, the power-3 closure performs quite well in this region. The advantage of the maxent closure becomes more noticeable on, and above the diagonal, which is associated with aggregated patterns. The ability to predict correctly the




Fig. 11 Relative error of the maxent closure (a) and the power-3 closure (b). We see that the maxent closure performs better than the power-three closure for mildly aggregated patterns (lower left), but the Kirkwood closure outperforms the maxent in segregated patterns (lower right)




Fig. 12 Difference in relative error between the maxent and power-3 closures for various combinations of dispersal and mortality spatial scales. Values higher than zero indicate that the maxent closure outperforms the power-3 closure, whereas negative values are evidence of better precision of the power-3 closure

equilibrium density in this regime is nearly optimal; particularly when the two scales have similar magnitudes, even when both dispersal and mortality act over short ranges. The regions of the parameter space for which each of the two closures is relatively more useful are shown in Fig. 12, which displays the difference in relative error between the two closures $\Delta E = err_{p3} - err_{maxent}$. Positive values of ΔE indicate that the error in the power-3 closure is larger than the maxent closure, and vice versa for negative

values of ΔE . As discussed above the largest improvement of the maxent closure around to the region where the two scales are of similar magnitude.

6 Discussion

The results of this research resonate with previous work (Bolker and Pacala 1997; Law et al. 2003; Ovaskainen and Cornell 2006a) that demonstrates that the analysis of stochastic, locally regulated, individual-based models of population dynamics in continuous space is feasible (Ovaskainen and Cornell 2006a; Bolker and Pacala 1997; Law et al. 2003). The numerical implementation of the maxent closure is computationally more expensive (about twice as much) than existing closure methods, but is nonetheless faster than resorting to direct simulation of the point process; if one is willing to approximate, the simplified closure based solely on the exponential correction (48) is substantially simpler to implement, and produces very small errors in comparison with the full maxent closure. Although a number of moment closures have been proposed in the literature, some using heuristic arguments, and others based on constrained entropy maximisation, very few, if any have a criterion of validity, with the exception of Ovaskainen and Cornell (2006a) who were able to derive a series expansion for the mean density of a spatially explicit metapopulation problem, and show rigorously that their approximation to the mean density is exact in the limit of long range interactions. The principal benefit of the maxent method lies in the fact that the normalisation constraint used to find the domain for the correction terms fails to find a non-trivial root when the closure assumptions are not met. This situation occurs when higher order terms are required in order to fully capture the dynamics, or when correlations extend over a range that goes beyond the window of observation. This property constitutes a validation check, not present in other proposed closure schemes.

Although the power-3 or Kirkwood closure had previously been derived from maximum entropy arguments (Singer 2004) (but using a different set of constraints and a different definition of the entropy functional), the correction terms presented here are new, and extend the probabilistic interpretation of the Kirkwood closure to situations where there is a region of irreducible triplet correlations. These correction terms introduce significant improvements in the agreement between the simulations of the stochastic process (for mildly aggregated patterns) and its deterministic approximation by the product density hierarchy. It remains to be seen how the maxent closure behaves for other functional forms of the interaction kernels, particularly for those that have tails that decay algebraically (i.e. power laws) instead of exponential. Another area of further work would be related to changes in the value of the non-spatial carrying capacity K . For higher densities, spatial effects become less important.

Since the derivation of the method does not depend on the details of the model, but only on that its equilibrium distribution is of maximum *locational* entropy with moment constraints, the maxent closure may be useful beyond spatial ecology where unclosed hierarchies for particle distribution functions are also commonly found, for instance in the statistical mechanics of fluids where the Kirkwood closure was first

introduced (Singer 2004), or in problems where the organisms move in space (Birch and Young 2006; Flierl et al. 1999; Young et al. 2001), provided that the correlation functions in those models are stationary in both space and time. A limitation of the method is its poor ability to predict the transient. This is to be expected, since maximum entropy is a meaningful property of the *equilibrium* distribution only when detailed balance is satisfied (Van Kampen 2001; Gardiner 1985; Jaynes 1957) and the transitions due to fecundity and dispersal events coincide with mortality. Other areas of current and future work include the generalisation of the moment hierarchy and the maxent closure to an arbitrary order of truncation, extensions to *marked* spatial point processes for populations with both spatial and size structure.

Acknowledgments M.R acknowledges the support granted by the International Institute for Applied Systems Analysis (IIASA) to participate in the Young Scientist Summer Program where part of this research was conducted during the summer of 2004. M.R. is grateful to Richard Law who suggested to work on this problem and generously shared his time and insights, Kenneth Lindsay who kindly shared his probabilistic and simulation expertise, and fruitful discussions with Jonathan Dushoff. The authors are also grateful with Benjamin Bolker, David Murrell and David Grey who helped with details on the simulations of the point processes and generously provided access to code and manuscripts. The support of Simon A. Levin and Ioannis G. Kevrekidis is gratefully acknowledged. The authors are grateful to an anonymous reviewer for insightful comments and suggestions.

Appendix: Derivation of moment equations

In order to derive the equation for $m_1(t)$, we start by fixing a small region of observation dx_1 (so that the count inside dx_1 , $N(dx_1)$ is either 0 or 1) and write a Master equation for the probabilities of change in the count $\Delta N_{\delta t}(dx_1)$ during a small time interval δt , defined as

$$\Delta N_{\delta t}(dx_1) = N_{t+\delta t}(dx_1) - N_t(dx_1).$$

These come from the birth and death transitions. Births are given by the probability that the count $N(dx_1)$ increases by one in δt due to a birth in dx_1

$$N \mapsto N + 1,$$

This probability is controlled by the fecundity rate and the dispersal kernel,

$$\begin{aligned} f(x_1|\varphi_t) &= \mathbb{P}\{\text{one birth in } (dx_1) \text{ during } (t, t + \delta t) \mid \varphi_t(X)\}. \\ &= \left[b \sum_{x_n \in \varphi_t} B(x_1 - x_n) N_t(dx_n) \ell(dx_1) \right] \delta t + o(\delta t), \end{aligned} \quad (55)$$

where b is the birth rate, $B(\xi)$ is the dispersal kernel, φ_t is the configuration of points at time t and $\ell(A)$ is the area of the 2-dimensional domain A . For the death of the individual in dx_1 , we have the transition

$$N \mapsto N - 1,$$

controlled by

$$\begin{aligned} \mu(x_1|\varphi_t) &= \mathbb{P}\{\text{death of individual } x_1 \text{ during } (t, t + \delta t) \mid \varphi_t(X)\} \\ &= N_t(dx_1) \left[d + d_N \sum_{x_n \in \varphi_t} W(x_1 - x_n) (N_t(dx_n) - \delta_{x_1}(dx_n)) \right] \delta t + o(\delta t), \end{aligned} \tag{56}$$

where d and d_N are positive constants defined in Sect. 2, the density-independent, and density-dependent contributions to the mortality and $W(\xi)$ is the mortality kernel). This probability is conditional on there being an individual in dx_1 . The change in the count $\Delta N_{\delta t}(x_1)$ is then given by both contributions

$$\Delta N_{\delta t}(dx_1) = f(x_1|\varphi_t) - \mu(x_1|\varphi_t)$$

so

$$\begin{aligned} \Delta N_{\delta t}(dx_1) &= \left[b \sum_{x_n \in \varphi_t} B(x_1 - x_n) N_t(dx_n) \ell(dx_1) \right. \\ &\quad \left. - N_t(dx_1) \left(d + d_N \sum_{x_n \in \varphi_t} W(x_1 - x_n) (N_t(dx_n) - \delta_{x_1}(dx_n)) \right) \right] \delta t. \end{aligned} \tag{57}$$

Taking expectations (ensemble averaging) on both sides and dividing by the duration of a small time interval δt yields

$$\begin{aligned} \frac{\mathbb{E}\{\Delta N_{\delta t}(dx_1)\}}{\delta t} &= b \sum_{x_n \in \varphi_t} B(x_1 - x_n) \mathbb{E}\{N_t(dx_n)\} \ell(dx_1) \\ &\quad - \mathbb{E}\left\{ N_t(dx_1) \left(d + d_N \sum_{x_n \in \varphi_t} W(x_1 - x_n) (N_t(dx_n) - \delta_{x_1}(dx_n)) \right) \right\}. \end{aligned}$$

after rearranging the second term, dividing both sides by $\ell(dx_1)$ and multiplying the second sum by $\ell(dx_n)/\ell(dx_n)$ we get

$$\begin{aligned} \frac{\mathbb{E}\{\Delta N_{\delta t}(dx_1)\}}{\ell(dx_1) \delta t} &= b \frac{\mathbb{E}\{N_t(dx_n)\}}{\ell(dx_1)} \sum_{x_n \in \varphi_t} B(x_1 - x_n) \ell(dx_1) - d \frac{\mathbb{E}\{N_t(dx_1)\}}{\ell(dx_1)} \\ &\quad - d_N \sum_{x_n \in \varphi_t} W(x_1 - x_n) \frac{\mathbb{E}\{N_t(dx_1) (N_t(dx_n) - \delta_{x_1}(dx_n))\}}{\ell(dx_1) \ell(dx_n)} \ell(dx_n). \end{aligned}$$

taking the limits as $\ell(dx_1)$ and $\ell(dx_n)$ go to zero, and using definition of the product density (16)

$$\begin{aligned} \frac{\Delta m_1(x_1, t)}{\delta t} &= b m_1(x_1, t) \int_{\mathfrak{R}^2} B(x_1 - x_n) dx_1 - d m_1(x_1, t) \\ &\quad - d_N \int_{\mathfrak{R}^2} W(x_1 - x_n) m_2(x_1, x_n, t) dx_n. \end{aligned}$$

since the process is spatially stationary by construction and exploiting the fact that the dispersal kernel integrates to unity, yields

$$\frac{\Delta m_1(t)}{\delta t} = b m_1(t) - d m_1(t) - d_N \int_{\mathfrak{R}^2} W(\xi_1) m_2(\xi_1, t) d\xi_1,$$

finally, after taking the limit as $\delta t \rightarrow 0$ we get,

$$\frac{d}{dt} m_1(t) = b m_1(t) - d m_1(t) - d_N \int_{\mathfrak{R}^2} W(\xi_1) m_2(\xi_1, t) d\xi_1. \tag{58}$$

On setting $r = b - d$, we get the generalisation of the logistic equation to the spatial case obtained by Law and Dieckmann (2000) and Law et al. (2003), but derived explicitly in terms of product densities,

$$\frac{d}{dt} m_1(t) = r m_1(t) - d_N \int_{\mathfrak{R}^2} W(\xi_1) m_2(\xi_1, t) d\xi_1. \tag{59}$$

Since m_2 is unknown, we need an additional evolution equation for this object. We follow a similar procedure to that used for the mean density, but considering the expected change of the *product* of the counts in two observation regions dx_1 and dx_2 . This requires the consideration of how pairs of points are created and destroyed as individuals disperse and die. There are three possible ways in which changes to occur. The first is to fix the count $N_t(dx_1)$ and allow only $N_t(dx_2)$ to change. The second is the reverse situation, fixing $N_t(dx_2)$ and allowing only $N_t(dx_1)$ to change. The third is when *both* $N_t(dx_1)$ and $N_t(dx_2)$ change in a small time interval. We have that

$$\begin{aligned} \Delta[N_t(dx_1) (N_t(dx_2) - \delta_{x_1}(dx_2))] &= N_t(dx_1) \Delta(N_t(dx_2) - \delta_{x_1}(dx_2)) \\ &\quad + (N_t(dx_2) - \delta_{x_1}(dx_2)) \Delta N_t(dx_1) \\ &\quad + \Delta N_t(dx_1) \Delta(N_t(dx_2) - \delta_{x_1}(dx_2)) \end{aligned} \tag{60}$$

where the Dirac delta distribution is used to remove self-pairs. The following derivation for the second order product densities is based on the symmetry in the probabilities of a birth or a death event occurring at both extremes of the distance vector linking x_1

and x_2 . We also assume that a *simultaneous* change in both $N_t(dx_2)$ and $N_t(dx_1)$ is negligible

$$\mathbb{P}[\Delta N_t(dx_1)\Delta(N_t(dx_2) - \delta_{x_1}(dx_2))] = o(\delta t)$$

and thus the transitions of second order can be written as

$$\Delta[N_t(dx_1) (N_t(dx_2) - \delta_{x_1}(dx_2))] = 2\Delta N_t(dx_1)(N_t(dx_2) - \delta_{x_1}(dx_2)). \tag{61}$$

Since we already have an expression for $\Delta N_t(dx_1)$, given by (57), (61) becomes

$$\begin{aligned} &\Delta[N_t(dx_1) (N_t(dx_2) - \delta_{x_1}(dx_2))] \\ &= 2 \cdot (N_t(dx_2) - \delta_{x_1}(dx_2)) \left[b \sum_{x_n \in \varphi_t} B(x_1 - x_n) N_t(dx_n) \ell(dx_1) \right. \\ &\quad \left. - N_t(dx_1) \left(d + d_N \sum_{x_n \in \varphi_t} W(x_1 - x_n)(N_t(dx_n) - \delta_{x_1}(dx_n)) \right) \right] \delta t. \end{aligned}$$

Taking expectations, and dividing by both sides by δt gives

$$\begin{aligned} \frac{\Delta[N_t(dx_1) (N_t(dx_2) - \delta_{x_1}(dx_2))]}{2 \delta t} &= b \sum_{x_n \in \varphi_t} B(x_1 - x_n) \mathbb{E} \{ N_t(dx_n)(N_t(dx_2) \\ &\quad - \delta_{x_1}(dx_2)) \} \ell(dx_1) \\ &\quad - d \mathbb{E} \{ N_t(dx_1)(N_t(dx_2) - \delta_{x_1}(dx_2)) \} \\ &\quad - d_N \sum_{x_n \in \varphi_t} W(x_1 - x_n) \mathbb{E} \{ N_t(dx_1)(N_t(dx_n) \\ &\quad - \delta_{x_1}(dx_n))(N_t(dx_2) - \delta_{x_1}(dx_2)) \}. \end{aligned}$$

After dividing by $\ell(dx_1)$ and $\ell(dx_2)$, using the definition of product densities (16) and taking the continuum limit in both space and time, one arrives at the evolution equation for the second order product density

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} m_2(\xi_1, t) &= b \int_{\mathfrak{N}^2} B(\xi_2) m_2(\xi_1 - \xi_2, t) d\xi_2 + b B(\xi_1) m_1(t) - d m_2(\xi_1, t) \\ &\quad - d_N W(\xi_1) m_2(\xi_1, t) - d_N \int_{\mathfrak{N}^2} W(\xi_2) m_3(\xi_1, \xi_2, t) d\xi_2, \tag{62} \end{aligned}$$

where we see the dependence on the *third* order product density in the last integral.

References

Birch DA, Young WR (2006) A master equation for a spatial population model with pair interactions. *Theor Popul Biol* 70(1):26–42

- Blaszczyszyn B (1995) Factorial moment expansion for stochastic systems. *Stoch Process Appl* 56:321–335
- Blath J, Etheridge A, Meredith M (2007) Coexistence in locally regulated competing populations and survival of branching annihilating random walk. *Ann Appl Probab* 17(5/6):1474–1507
- Bolker B, Pacala SW (1997) Using moment equations to understand stochastically driven spatial pattern formation in ecological systems. *Theor Popul Biol* 52:179–197
- Bolker B, Pacala SW (1999) Spatial moment equations for plant competition: understanding spatial strategies and the advantages of short dispersal. *Am Nat* 153(6):575–602
- Bolker B, Pacala SW, Neuhauser C (2003) Spatial dynamics in model plant communities: what do we really know. *Am Nat* 162(2):135–148
- Bolker B, Pacala SW, Levin SA (2000) Moment methods for ecological processes in continuous space. In: Dieckmann U, Law R, Metz JA (eds) *The geometry of ecological interactions*. Cambridge studies in adaptive dynamics, vol 1. Cambridge University Press, London, pp 388–411
- Borgogno F, D’Odorico P, Laio F, Ridolfi L (2009) Mathematical models of vegetation pattern formation in ecohydrology. *Rev Geophys* 47
- Brooker RW, Maestre FT, Callaway RM, Lortie CL, Cavieres LA, Kunstler G, Liancourt P, Tielboerger K, Travis JMJ, Anthelme F, Armas C, Coll L, Corcket E, Delzon S, Forey E, Kikvidze Z, Olofsson J, Pugnaire F, Quiroz CL, Saccone P, Schifffers K, Seifan M, Touzard B, Michalet R (2008) Facilitation in plant communities: the past, the present, and the future. *J Ecol* 96(1):18–34
- Condit R, Ashton PS, Baker P, Bunyavejchewin S, Gunatilleke S, Gunatilleke N, Hubell SP, Foster R, Itoh A, LaFrankie JV, Seng Lee H, Losos E, Manokaran N, Sukumar R, Yamakura T (2000) Spatial patterns in the distribution of tropical tree species. *Science* 288:1414–1418
- Cressie NA (1991) *Statistics for spatial data*. Wiley, New York
- Dale MR (1999) *Spatial pattern analysis in plant ecology*. In: Cambridge studies in ecology. Cambridge University Press, London
- Daley D, Vere-Jones D (1988) *An introduction to the theory of point processes*. In: Springer series in statistics. Springer, Berlin
- Daley D, Vere-Jones D (2003) *An introduction to the theory of point processes*. Volume I: elementary theory and methods. In: Probability and its applications, 2nd edn. Springer, Berlin
- Daley D, Vere-Jones D (2008) *An introduction to the theory of point processes*. Volume II: general theory and structure. In: Probability and its applications, 2nd edn. Springer, Berlin
- Daley DJ, Vere-Jones D (2004) Scoring probability forecasts for point processes: the entropy score and information gain. *J Appl Probab* 41A:297–312
- Damgaard C (2007) On the hyperbolic competition model: a comment to Weigelt et al. (2007). *J Ecol* 95(4):599–600
- DeAngelis DL, Mooij WM (2005) Individual-based modeling of ecological and evolutionary processes. *Annu Rev Ecol Evol Syst* 36:147–168
- Dieckmann U, Law R (2000) Relaxation projections and the method of moments. In: Dieckmann U, Law R, Metz JA (eds) *The Geometry of Ecological Interactions*. Cambridge studies in adaptive dynamics, vol 1. Cambridge University Press, London, pp 412–455
- Dieckmann U, Law R, Metz JA (eds) (2000) *The geometry of ecological interactions*. In: Cambridge studies in adaptive dynamics, vol 1. Cambridge University Press, London
- Diggle PJ (1983) *Statistical analysis of spatial point patterns*. Academic Press, London
- Dormann C, Roxburgh S (2005) Experimental evidence rejects pairwise modelling approach to coexistence in plant communities. *Proc R Soc B Biol Sci* 272(1569):1279–1285
- Durrett R (1999) Stochastic spatial models. In: Capasso V, Dieckmann O (eds) *Mathematics inspired by biology*. Lecture notes in mathematics, vol 1714. Springer, Berlin, pp 39–94
- Durrett R, Levin SA (1994) Stochastic spatial models: a user’s guide to ecological applications. *Philos Trans R Soc Lond B* 343:329–350
- Etheridge AM (2004) Survival and extinction in a locally regulated population. *Ann Appl Probab* 14(1):188–214
- Filipe J, Gibson G (1998) Studying and approximating spatio-temporal models for epidemic spread and control. *Philos Trans R Soc Lond Ser B Biol Sci* 353(1378):2153–2162
- Filipe J, Gibson G (2001) Comparing approximations to spatio-temporal models for epidemics with local spread. *Bull Math Biol* 63(4):603–624
- Filipe J, Maule M (2003) Analytical methods for predicting the behaviour of population models with general spatial interactions. *Math Biosci* 183(1):15–35
- Filipe J, Maule M, Gilligan C (2004) On ‘Analytical models for the patchy spread of plant disease’. *Bull Math Biol* 66(5):1027–1037

- Flierl H et al (1999) From individuals to aggregations: the interplay between behavior and physics. *J Theor Biol* 196:397–454
- Fournier N, Méléard S (2004) A microscopic probabilistic description of a locally regulated population and macroscopic approximations. *Ann Appl Probab* 14(4):1880–1919
- Gardiner C (1985) *Handbook of stochastic methods*, 2nd edn. Springer, Berlin
- Gillespie D (1976) General method for numerically simulating stochastic time evolution of coupled chemical reactions. *J Comput Phys* 22:403–434
- Gratzer G, Canham C, Dieckmann U, Fisher A, Iwasa Y, Law R, Lexer MJ, Sandmann H, Spies TA, Splechtna BE, Szwagryzk J (2004) Spatio-temporal development of forests-current trends in field methods and models. *Oikos* 107(1):3–15
- Grimm V (1999) Ten years of individual-based modelling in ecology: what have we learned and what could we learn in the future?. *Ecol Modell* 115:129–148
- Gross K (2008) Positive interactions among competitors can produce species-rich communities. *Ecol Lett* 11:929–936
- Grouba V, Zorin A, Sevastianov L (2004) The superposition approximation: a critical review. *Int J Mod Phys B* 18(1):1–44
- Guan Y (2007a) A composite likelihood cross-validation approach in selecting the bandwidth for the estimation of the pair correlation function. *Scand J Stat* 34(2):336–346
- Guan Y (2007b) A least-squares cross-validation bandwidth selection approach in pair correlation function estimations. *Stat Probab Lett* 77:1722–1729
- Hiebeler D (1997) Stochastic spatial models: from simulations to mean field and local structure approximations. *J Theor Biol* 187(3):307–319
- Hiebeler D (2006) Moment equations and dynamics of a household SIS epidemiological model. *Bull Math Biol* 68(6):1315–1333
- Hillen T (2004) On the L^2 -moment closure of transport equations: the cattaneo approximation. *Discret Continuous Dyn Syst Ser B* 4(4):961–982
- Iwasa Y (2000) Lattice models and pair approximations in ecology. In: Dieckmann U, Law R, Metz JA (eds) *The geometry of ecological interactions*. Cambridge studies in adaptive dynamics, vol 1. Cambridge University Press, London, pp 227–251
- Janossy L (1950) On the absorption of a nucleon cascade. *Proc R Ir Acad Sci Sect A* 53:181–188
- Jaynes ET (1957) Information theory and statistical mechanics. *Phys Rev* 106(4):620–630
- Jaynes ET (1982) On the rationale of maximum entropy methods. *Proc IEEE* 70(9):939–952
- Keeling M (1999) The effects of local spatial structure on epidemiological invasions. *Proc R Soc Lond Ser B Biol Sci* 266(1421):859–867
- Keeling M, Rand D, Morris A (1997) Correlation models for childhood epidemics. *Proc R Soc Lond Ser B Biol Sci* 264(1385):1149–1156
- Keeling M, Wilson H, Pacala S (2000) Reinterpreting space, time lags, and functional responses in ecological models. *Science* 290(5497):1758–1761
- Keeling MJ, Ross JV (2009) Efficient methods for studying stochastic disease and population dynamics. *Theor Popul Biol* 75(2-3):133–141
- Khinchin AI (1957) *Mathematical foundations of information theory*. Dover, New York
- Kirkwood JG, Boggs EM (1942) The radial distribution function in liquids. *J Chem Phys* 10:394–402
- Krishnarajah I, Cook A, Marion G, Gibson G (2005) Novel moment closure approximations in stochastic epidemics. *Bull Math Biol* 67(4):855–873
- Law R, Dieckmann U (2000) Moment approximations of individual-based models. In: Dieckmann U, Law R, Metz JA (eds) *The geometry of ecological interactions*. Cambridge studies in adaptive dynamics, vol 1. Cambridge University Press, London, pp 252–270
- Law R, Murrell DJ, Dieckmann U (2003) Population growth in space and time: spatial logistic equations. *Ecology* 84(1):252–262
- Levin SA (1994) Patchiness in marine and terrestrial systems: from individuals to populations. *Philos Trans R Soc Lond Ser B* 3:99–103
- Lewis M, Pacala S (2000) Modeling and analysis of stochastic invasion processes. *J Math Biol* 41(5):387–429
- McFadden J (1965) The entropy of a point process. *J Soc Ind Appl Math* 12(4):988–994
- Meeron E (1957) Series expansion of distribution functions in multicomponent fluid systems. *J Chem Phys* 27(6):1238–1246
- Moorcroft PR, Pacala SW (2001) A method for scaling vegetation dynamics: the ecosystem demography model. *Ecol Monogr* 71(4):557–586

- Murray J (1992) *Mathematical Biology*. In: Springer series in applied mathematics, vol 85. Springer, Berlin
- Murray JD (1993) *Mathematical biology*. In: Biomathematics, vol 19. Springer, Berlin
- Murrell D, Dieckman U, Law R (2004) On moment closures for population dynamics in continuous space. *J Theor Biol* 229(3):421–432
- Othmer H, Dunbar S, Alt W (1988) Models of dispersal in biological systems. *J Math Biol* 26:263–298
- Ovaskainen O, Cornell SJ (2006a) Asymptotically exact analysis of stochastic metapopulation dynamics with explicit spatial structure. *Theor Popul Biol* 69:13–33
- Ovaskainen O, Cornell SJ (2006b) Space and stochasticity in population dynamics. *Proc Natl Acad Sci* 103(34):12,781–12,786
- Pacala SW, Levin SA (1997) Biologically generated spatial pattern and the coexistence of competing species. In: Tilman D, Kareiva P (eds) *Spatial ecology: the role of space in populations and interspecific interactions*. Monographs in population biology, vol. 30, chap. 9. Princeton University Press, New Jersey, pp 204–232
- Pascual M, Levin SA (1999) From individuals to population densities: searching for the intermediate scale of nontrivial determinism. *Ecology* 80(7):2225–2236
- Pascual M, Mazzega P, Levin S (2001) Oscillatory dynamics and spatial scale: the role of noise and unresolved pattern. *Ecology* 82(8):2357–2369
- Pascual M, Roy M, Franc A (2002) Simple temporal models for ecological systems with complex spatial patterns. *Ecol Lett* 5(3):412–419
- Pearl R, Reed L (1920) On the rate of growth of the population of the United States since 1790 and its mathematical representation. *Proc Natl Acad Sci* 6:275–288
- Picard N, Franc A (2001) Aggregation of an individual-based space-dependent model of forest dynamics into distribution-based and space-independent models. *Ecol Model* 145(1):69–84
- Purves DW, Law R (2002) Experimental derivation of functions relating growth of *Arabidopsis thaliana* to neighbour size and distance. *J Ecol* 90:882–894
- Raghieb Moreno M (2006) Point processes in spatial ecology. Ph.D. thesis, Department of Mathematics, University of Glasgow
- Renshaw E (1991) *Modelling biological populations in space and time*. In: Cambridge studies in mathematical biology, vol 11. Cambridge University Press, London
- Salpeter E (1958) On mayers theory of cluster expansion. *Ann Phys* 5(3):183–223
- Satō K, Iwasa Y (2000) Pair approximations for lattice-based ecological models. In: Dieckmann U, Law R, Metz JA (eds) *The geometry of ecological interactions*. Cambridge studies in adaptive dynamics, vol 1. Cambridge University Press, London, pp 341–358
- Scanlon TM, Caylor KK, Levin SA, Rodriguez-Iturbe I (2007) Positive feedbacks promote power-law clustering of Kalahari vegetation. *Nature* 449(7159):209–212
- Schneider MK, Law R, Illian JB (2006) Quantification of neighbourhood-dependent plant growth by bayesian hierarchical modelling. *J Ecol* 94(2):310–321
- Scott DW (1992) *Multivariate density estimation: theory, practice, and visualization*. In: Wiley series in probability and statistics. Wiley, New York
- Sese L (2005) Triplet correlations in the quantum hard-sphere fluid. *J Chem Phys* 123(10)
- Shannon CE, Weaver W (1949) *The mathematical theory of communication*. University of Illinois Press, Champaign
- Silvertown J, Doust JL (1993) *Introduction to plant population biology*. Blackwell, Oxford
- Silvertown J, Wilson JB (2000) Spatial interactions among grassland plant populations. In: Dieckmann U, Law R, Metz JA (eds) *The geometry of ecological interactions*. Cambridge studies in adaptive dynamics, vol 1. Cambridge University Press, London, pp 28–46
- Singer A (2004) Maximum entropy formulation of the Kirkwood superposition approximation. *J Chem Phys* 121(8):3657–3666
- Stoll P, Weiner J (2000) A neighborhood view of interactions among individual plants. In: Dieckmann U, Law R, Metz JA (eds) *The geometry of ecological interactions*. Cambridge studies in adaptive dynamics, vol 1. Cambridge University Press, London, pp 11–27
- Stoyan D, Helga S (1994) *Fractals, random shapes and point fields*. Wiley, New York
- Stoyan D, Kendall WS, Mecke J (1995) *Stochastic geometry and its applications*. In: Wiley series in probability and statistics, 2nd edn. Wiley, New York
- Turnbull LA, Coomes DA, Purves DW, Rees M (2007) How spatial structure alters population and community dynamics in a natural plant community. *J Ecol* 95(1):79–89
- Van Kampen N (2001) *Stochastic processes in physics and chemistry*. North-Holland, Amsterdam

- Verhulst P (1838) Notice sur la loi que la population suit dand son accroissement. *Corres Math Phys* 10:113–121
- Weigelt A, Schumacher J, Walther T, Bartelheimer M, Steinlein T, Beyschlag W (2007) Identifying mechanisms of competition in multi-species communities. *J Ecol* 95(1):53–64
- Wilson HB, Keeling MJ (2000) Spatial scales and low-dimensional deterministic dynamics. In: Dieckmann U, Law R, Metz JA (eds) *The geometry of ecological interactions*. Cambridge studies in adaptive dynamics, vol 1. Cambridge University Press, London, pp 209–226
- Young WR, Roberts AJ, Stuhne G (2001) Reproductive pair correlations and the clustering of organisms. *Nature* 412:328–331